

Frequent Occurrence of Sports Injuries amid Female Athletes during Practice and Competition Events

Charan Singh*

Assistant Professor, Department of Physical Education, Baba Balraj, Panjab University Constituent College, Balachaur

Abstract – Numerous epidemiological investigations in montages and professional athletes in both genders uncover that ankle and knee are the anatomical destinations that are regularly injured. Sprains, strains and breaks are the normal sort of injuries. The reason for the examination is to analyze the study of disease transmission of practice and competition injury in young men and young ladies sports.

Keywords: Sports Injuries, Female Athletes, Practice Session, Competition Events

INTRODUCTION

The sports injuries are isolated into acute and abuse mishaps. Acute mischances are distressing injuries which incorporate breaks disengagement, ligament sprains and muscle tears, and abuse injuries, comprehensive of stress crack compartment disorders and tendinopathies. This is totally a result of the duplicated load set upon musculoskeletal shape by methods for the increased schooling needs of current day amusement and increased acknowledgment of ingenuity exercises.

The numerous legends that ladies are the "weaker sex" and in this way need to best enjoys positive proper sports exercises comprising of tennis, playing golf and maybe experiencing and horseback utilizing, with specific obstructions, no genuine examination has been embraced with regards to the real physical capacities of the ladies competitor till most recent years.

Females have end up being progressively more stressed in conveying side interest in the course of recent decades. Female athletes by and large have a tendency to extend comparable mischief issues to their male partners. In any case, some of zones are of specific subject to women.

Anyway young ladies had their partner to the authentic Olympics with the Games of Hera, These games in which women hustled, over the space of around 500 toes performed almost no notoriety in verifiable Greek human advancement. Aristotle summed up the noteworthy Greek's state of mind to the ladies while he composed the essayist of nature

gave fellow power of casing and mettle of brain to empower him to confront outstanding inconvenience and to young ladies gave a frail and delicate constitution executed by methods for a home grown delicateness and lowliness which sound her for partisan presence.

The resurrection of present day Olympics occurred inside the Victorian time when young ladies had been showed up as powerless, delicate and latent. As a final product of this mentality no young lady exercises folks held inside the primary present day Olympiad in 1896 and best two sports exercises, tennis and golf, had lady contenders inside the 1900 Olympics in Paris.

Since those early Olympics there was a bit by bit expanding participation of ladies inside the Olympic Games in each mid-year and frigid climate computer games. The main team activity conveyed for women who had been presented simultaneously for men and young ladies progressed toward becoming volleyball in 1964. Proof that lady association in the Olympic Games had 'come to age' wind up demonstrated in Los Angeles in 1984 while the most discussed exercises of the computer games transformed into encounter among Many Decker-Slaney and Zola Budd in 3000 m. Anyway social and non-mainstream confinement women that the huge amount of nations don't allow female participation in game.

The expansion in female participation in game isn't just restricted to contribution in the Olympic Games. There has been a sensational development inside the assortment of young lady practicing and contending in game. It is presently an ordinary sight in most extreme western global areas to look women running,

taking strolls and going to senobc directions notwithstanding contending in character and group activities exercises. While a couple of female competitor and female sports exercises have accomplished an observably over the top profile and money related reestablished, most lady game isn't bolstered by method for media and ability supports. The gigantic livelihoods of driving female tennis players are the exemption as a substitute the main run the show.

There are as yet solid confinements in our general public that keep ladies from transforming into stressed in sport a couple of individuals from society sees sports exercises sports women as 'butch' or unfeminine!, solid social for high accomplishment and most satisfaction in exercise.

Guardians have a basic task to carry out in digging in all children no longer only children to be engaged with sport. Female stressed in physical intrigue tend to have better confidence other those not stressed. Being amazing at game positions especially as a trait for more youthful men yet it isn't positioned moderately as a trademark for youthful guys anyway it isn't positioned phenomenally as a quality for young ladies, either by means of women or by means of men. This outlook may be a prime thing in high drop-out from female amusement round adolescence.

Much advancement has been made in women game inside the previous 3 numerous years anyway comparably endeavors a required to move young ladies to assimilate and hold to separated. The most critical brief era reason should be to blast the overall low quantities of young lady in training and authoritative jobs. This incorporates the wide assortment of lady sports therapeutic medication specialists.

RESEARCH OBJECTIVES

Our goal is to look at of practice and consummation injuries in 5 young lady's sports (softball volleyball, container ball, aerobatic and running). For these events the particular points are the accompanying.

- To study the distinctive sports injuries of female athletes managed in practice.
- To examine the different sports injuries of female athletes supported in competition.
- To compare the sports injuries of female competitor in practice and competition.

REVIEW OF LITERATURE

The larger part of research executed have had a tendency to view handiest the physiological segment of molding programs for ladies through assorted methods of exercise. They have estimated strong tissues, persistence bends, fundamental ability and

distinctive elements and they have likewise analyzed gynecological issues of lady competitor.

All the sport requests appropriate real wellbeing comprising of harmony, control, vitality and adaptability. Amid the positive computer games the players take off, turn, keep running ahead and in reverse and exchange course customarily. These sports with the strain to be triumphant more noteworthy paints and to acknowledge as conceivable less put radical mechanical powers in musculoskeletal framework which commonly injured. More over the play area plays a choice capacity inside the example of injury.

Rechel et al, 2008, completed the inspect on an epidmioligical difference of high school sports mishaps maintained in practice and competition their take a gander at well-knownshows that rates and examples of over the top personnel sports injuries contrasted among practice and competition.

Haycock et al 201976 concentrated the defenselessness of women Athletes to hurt in USA. They found that harm maintained by utilizing female competitor, relegated by means of those identified with exceptional natural shape, were basically no longer extraordinary them the ones of colleagues nor are pleasantly prepared woman athletes more noteworthy powerless against mishaps.

A potential see on b-ball injuries among male and lady athletes progressed toward becoming directed by means of Stergiolelas et. A; 2007. They look at that the cost of mischances as per male competitor in accordance with a year changed into 0. Seventy two while for young ladies the comparing rate for zero. Fifty six. Male competitor her circle abuse injuries than females ($p < \text{zero}.02$), in the meantime as female had pass harm in lumbar spine ($p < 0.001$). Male had additional respected sprain in average insurance ligament ($p < \text{zero}.05$) even as young lady had stream sprain in front cruciate ligament ($p < \text{zero}.1/2$).

A planned inspect of abuse knee injuries changed into did by means of Devan et al 2004, among lady Athletes with muscle umbalances and based variations from the norm their results shows that Ten abuse knee injuries iliotibial band grating disorder = S, patellar tendintis = 3, patellofemoral syndrome=1, pes onserine lendinitis=1) came to pass for in gathetes. The H:Q proportion under the regular nouge at three hundred% ($P=0.047$) wound up identified with abuse knee mishaps, as turn into the nearness of genu recon weight ($p=0.004$) moreover.

Haycock et al 1976 concentrated the vulnerability of women. Athletes to hurt in USA. They found that injury managed by methods for woman competitor, doled out by method for those related with unique natural structure, have been basically not unmistakable them the ones of men, nor are

legitimately talented young lady athletes more noteworthy in danger of mischances.

A forthcoming take a gander at on b-ball mishaps among male and lady athletes changed into performed through Stergioulelas et-al 2007.

They investigate that the expense of injuries per artistic creations competitor with regards to a year wound up zero. Seventy two even concerning lady lies the relating charge for zero. Fifty six Male competitor her more prominent over utilize mishaps than females ($p < 0.02$), in the meantime as female had pass injury in Lumbar spine ($p < 0.001$). Male had more prominent Kumar Sprain in average collateral ligament ($p < 0.05$) even as young lady had stream sprain in front cruciate ligament ($p < 0.1/2$) athletes squeezing diminish H:Q proportions at three hundred zero/s and genu recurvatum brought about additional over utilize knee mischances than athletes without these irregularities ($P = 0.001$)

MATERIAL AND METHODS

Justification of the problem: The motivation behind this examination is to record injuries and relative investigation of Sports injuries in female athletic managed in practice and competition. There has been a developing enthusiasm for ladies' sports from, softball, crate ball, volleyball, tumbling and track. As of late young ladies have begun to take an interest in more athletic events, which used to be overwhelmed by young men and young fellows. A couple of years prior ladies were considered to play a slower, more cautious style of sports. Nonetheless, today they are more aggressive. Ladies are endeavoring to get grant and professional contracts or win events that will bring them monetary benefit end the open door for underwriting by huge name organizations. This has made ladies play with more power and speed. Their more prominent participation end force of play has expanded the level of enthusiasm for wellbeing and restorative issues particular to female games. So the motivation behind why specialist pick this relative investigation on the grounds that there are restricted examination in Delhi and NCR district.

RESEARCH DESIGN

The descriptive type of survey method has been used.

Population: Here the population is the all-female athletes of Patiala city.

Sample: 200 female athletes were selected randomly from Patiala city.

Tools used: A self-made questionnaire was used for the present paper.

Subject: Volunteers were from different Physical Education Colleges in Patiala (Punjab). All athletes marked educated assent shapes. This number of female athletes picked was randomly selected amid the period June 2018 - August 2018. Mentors and players, after the determination, were educated in insight about the reason for study and the manner in which information would be gathered. The mentors were given a standard sheet from examiner. The information included: anatomical locale of injury and period however injury happened (amid practice or competition).

For assessment of injury rate: The quantity of female athletes that incorporated into the examination was separated by number of female competitor injured. The athletes were selected who were in practice or competition.

For this examination an injury could be characterized as a condition meeting the accompanying 3 criteria's:

- Occurred because of partaking in a composed college practice and fulfillment.
- Required therapeutic consideration by an AT or doctor and.
- Resulted in confinement of the student athletics participation for 1 day or more past the day of injury.

The competitor finished injury reports which incorporate competitor socioeconomics (E.g.- age, weight, year in school) and the conditions encompassing the injury occasion (E.g.- instrument, time in practice and competition, position played). Rate proportions were ascertained with 95% certainty interim. A $RR > 1.00$ recommends a hazard affiliation while as $RR < 1.00$ proposes a defensive affiliation.

$$RR = \frac{[(\text{No. of Competition Injuries} / \text{No. of Competition AEs}) \times 1000]}{[(\text{No. of Practice Injuries} / \text{No. of Practice AEs}) \times 1000]}$$

RESULTS & DISCUSSION




Result

Injuring Rate: In the year 2017– 2018, athletes partaking in 5 young ladies sports (volley ball b-ball, delicate ball, Gymnastic and Running) at a national agent test of 3 schools maintained 54 injuries (26 in practice and 28 in competition) In can be seen from the table 1 that these injuries happened amid 200 Athlete Exposure (144 in practice and 50 in competition An and S) bringing about an aggregate injury rate of 270 POC 10 An and S. The rate of injury per 10 An and S was higher in competition (0.5) than in practice ($RR = 237, 95\%$) In practice the most astounding rate of injury 10 An and S happened in gymnastic (0.27), trailed by running (0.24) and b-ball (0.147). In competition the higher rate of injury per 10




AEs happened in gymnastic. (0.8) trailed by delicate ball (0.636) and running (0.6). The given information of injury rate amid competition and practice has been appeared by the accompanying diagram:- 1 to 7.


Table -1: Practice & competition injury rates

Sports	Injuries		Exposure		Rate of Injury (Per 10 Athlete)	Rate Ratio
	No.	Percentage	No.	Percentage		
Overate	59		200		2.70	
Practice	26	48	144	72	0.180	2.73
Competition	28	52	56	28	0.5	
Volleyball	12		52		2.30	
Practice	4	33.3	32	61.5	0.125	0.05
Competition	8	66.6	20	38.4	0.40	
Basketball	11		48		2.29	
Practice	5	45.4	33	70.8	0.147	3.40
Competition	6	54.5	15	25.0	0.5	
Soft ball	10		36		2.77	
Practice	3	30.0	25	69.4	0.12	5.3
Competition	7	70.0	11	30.5	0.363	
Gymnastic	12		34		3.52	
Practice	8	66.6	29	85.2	0.27	2.96
Competition	4	33.3	5	14.7	0.8	
Running	09		30		0.3	
Practice	06	66.6	25	83.3	0.24	2.5
Competition	03	33.3	5	16.6	0.6	



Graph





Discussion:

I even have found and portrayed gorged practice in 5 women sports exercises the utilization of certainties gathered among an agent example of real schooling understudies in Patiala and saw that restriction events results in higher injury rates. With creating participation in these athletes, the final product offers AEs with an all-around coordinated assessment of activity and competition mishances, a basic advance toward making sense of focused, confirmation based indications for higher injury counteractive action among athletes.

A few different creators have found that injury cites specifically sports exercises together with volleyball, ball, acrobatic and walking had been exceptional in resistance. Higher cost of damage all through resistance might be a result of lifted play profundity, extended physical touch and stretched out attention to over the top hazard sports. In spite of the better injury rate throughout competition it wound up discovered that practically identical offer of harm was managed in competition (52%) and work out (forty eight%). Likewise steady with the impacts distributed beforehand, this finding is a direct result of the more noteworthy record of 5 spent in 15 practice contrasted and restriction.

CONCLUSION

Our forthcoming investigation in female athletes maintained in practice and competition vents uncovered that injury rate is higher amid competition vents and practice sessions. In 5 young ladies sports, we found that in gymnastic the injury rate is significantly higher amid competition than practice sessions as pursued by bin ball, volleyball, softball and running. Additionally studies might be led to discover conceivable causes contributing injuries among ladies athletes.

REFERENCES

Beynon B.D., Vacek P.M., Murphy D., Alosa D., Paller D. (2005). First-time inversion ankle ligament trauma: the incidence of injury. *Am J sports Med.* 2005;33 (10): pp. 1485-1491.

Centers for Disease Control and Prevention (CDC) (2006). Sports-related injuries among high school athletes- United States, 2005-06 school year. *MMWR Morb Mortal Wkly Rep.* 2006; 55 (38): pp. 1037-1040.

Emery C.A., Meeuwisse W.H., Hartmann S.E. (1996). Evaluation of risk factors for injury in adolescent soccer: implementation and validation of an injury surveillance system. *Am J Sports Med.* 1996; 24(5): pp. 684-687.

Field M., Collins M.W., Lovell M.R., Maroon J. (2003). Does age play a role in recovery from sports-related concussion? A comparison of high school and collegiate athletes. *J. Pediatr.* 2003;142 (5): pp. 546-553.

Gavrilides A. sotiropoulos, M. and Balto Ponles P. (2007). Amateur basketball injuries. A prospective study among male and female athletes. (*SAJSM Vol.21 No.4*)

Guskiewicz K.M., Weaver N.L., Padua D.A., Garrett W.E. (2000). Jr, Epidemiology of concussion in collegiate and high school football players. *Am J sports Med.* 2000; 28(5): pp. 643-650.

Halpern B., Thompson N., Curl W.W., Andrews J.R., Hunter S.C., Boring J.R. (1987). High school football injuries: identifying the risk factors. *Am J Sports Med.* 1987;15 (4): pp. 316-320.

Haycock E.E. Joam M.D., Gillette V. (1976). Susceptibility of women athletes to injury (*Journal of JAMA July Vol.236. No.2*)

Hinton R.Y., Lincoln A.E., Almquist J.L., Douguih W.A., Sharma K.M. (2005). Epidemiology of lacrosse injuries in high school-aged girls and boys: a 3-year prospective study. *Am J Sports med,* 2005; 33(9): pp. 1305-1314.

Inklaar H. (1994). Soccer injuries, I: incidence and severity. *Sports Med.* 1994;18(1): pp. 55-73.

Ireland M.L. (1999). Anterior cruciate ligament injury in female athletes: epidemiology. *J. Athl Train.* 1999;34 (2): pp. 150-154.

Knowles S.B., Marshall S.W., Bowling J.M., et. al. (2006). A prospective study of injury incidence among North Carolina high school athletes. *Am J Epidemiol.* 2006; 164 (12): pp. 1209-1221.

McCrea M., Hammeke T., Olsen G., Leo P., Guskiewicz K. (2004). Unreported concussion in high school football players: implications for prevention *Clin J Sport Med.* 2004;14(1): pp. 13-17.

- Messina D.F., Farney W.C., DeLee U.J.C. (1999). The incidence of injury in Texas high school basketball: a Prospective study among male and female athletes. *Am J Sports Med.* 1999; 27(3): pp. 294-299.
- Mueller F.O. (2001). Catastrophic head injuries in high school and collegiate sports. *J Athl Train.* 2001;36(3): pp. 312-315.
- Nation Federation of State High School Associations (2006). 255-06 High school athletics participation survey. http://nfhs.org/core/contentmanager/uploads/2005_06_NFHS_participations_survey.pdf. Accessed November 2006.
- National Federation of State High School Associations (2007). The case for high school activities. http://www/nfhs.org/web/2004/01/the_case_for_high_school_activities.aspx, Accessed February 6,2007.
- Pasque C.B., Hewett T.E. (2000). A prospective study of high school wrestling injuries. *AM J Sports Med.* 2000; 28(4): pp. 509-515.
- Powell J.W., Barber-Foss K.D. (1999). Injury patterns in selected high school sports: a review of the 1995-1997 seasons. *J Athl Train.* 1999; 34(3): pp. 277-284.
- Powell J.W., Barber-Foss K.D. (1999). Traumatic brain injury in high school athletes. *JAMA.* 1999; 282(10): pp. 937-944.
- Powell J.W. (2001). Cerebral concussion: causes, effects, and risks in sports. *J Athl Train,* 2001;36 (3): pp. 307-311.
- Ramirez M., Schaffer K.B., Shen H., Kashani S., Kraus J.F. (2006). Injuries to high school football athletes in California. *Am J Sports Med.* 2006;34(7): pp. 1147-1158.
- Reshel, J.A.; E.E; comstole. R.D. (2008). An epidemalologic comparison of high school sports enquiries sustained in preactice and competition.
- Schulz M.R., Marshall S.W., Mueller F.O., et. al. (2004). Incidence and risk factors for concussion in high school athletes, North Carolina, 1996-1999. *Am J Epidemiol.* 2004;160 (10): pp. 937-944.
- Shankar P.R., Fields S.K., Collins C.L., Dick R.W., Comstock R.D. (2007). epidemiology of high school and collegiate football injuries in the United States, 2005-2006. *Am J Sports Med.* 2007;35(8): pp. 1295-1303.
- Stergioulas A., Tripolirioli A., Kostopoulos N., Stuart M.J. (2005). Gridiron football injuries. *Med Sport Sci.* 2005; 49: pp. 62-85.
- Turbeille S.D., Cowan L.D., Owen W.L., Asal N.R., Anderson M.A. (2003). Risk factors for injury in high school football players. *Am J Sports Med.* 2003;31(6): pp. 974-980.
- Willer B., Kroetsch B., Darling S., Hutson A., Leddy J. (2005). Injury rates I house league select, and representative youth ice hockey. *Med Sci sports Exerc.* 2005; 37(10): pp. 1658-1663.
- Wroble R.R., Mysnyk M.C., Foster D.T., Albright J.P. (1986). Patterns of knee injuries in wrestling a six year study. *Am J Sports Med.* 1986;14(1): pp. 55-66.

Corresponding Author

Charan Singh*

Assistant Professor, Department of Physical Education, Baba Balraj, Panjab University Constituent College, Balachaur

E-Mail – scharan1979@gmail.com