

Efficacy of Physiotherapy Exercises after Elective Total Knee Arthroplasty

Abdullah S. Bin Hussein¹ Nezar Al Torairi² Shabana Khan³ Sharick Shamsi^{4*}

¹ Officer, Prince Sultan Military Medical City, Riyadh KSA

² Clinical Supervisor at Ortho Physiotherapy OPD, at Prince Sultan Military Medical City, Riyadh KSA

³ Physiotherapist at Prince Sultan Military Medical City, Riyadh KSA

⁴ Senior Physiotherapist at Prince Sultan Military Medical City, Riyadh KSA

Abstract – The fundamental point of this paper is to assess the adequacy of physiotherapy exercise after elective primary total knee arthroplasty in patients with osteoarthritis. A short introduction is given to comprehend this theme all the more plainly. A survey of different investigations is additionally given. Randomized controlled preliminaries were investigated on the off chance that they incorporated a physiotherapy exercise intercession contrasted and common or standard physiotherapy care, or looked at two sorts of exercise physiotherapy mediations meeting the survey criteria, after releasing from the hospital after elective primary total knee arthroplasty for osteoarthritis. We efficiently checked on randomized controlled preliminaries to decide the adequacy of physiotherapy exercise after release as far as improving capacity, personal satisfaction, strolling, the scope of movement in the knee joint, and muscle quality for patients with osteoarthritis after elective primary one-sided total knee arthroplasty.

Keywords: Physiotherapy, Osteoarthritis, Knee, Arthroplasty

-----X-----

1. INTRODUCTION

Osteoarthritis is the commonest reason for inability in more seasoned individuals, with agonizing knee osteoarthritis influencing 10% of individuals matured more than 55 in everywhere throughout the world. Over 80% of patients' experience restrictions in performing exercises of everyday living, for example, portability outside the home, family unit tasks, and work obligations. In 2005, patients with osteoarthritis represented at any rate 55,495 primary knee joint arthroplasties. As the length of the hospital remains after joint arthroplasty surgery has notably and quickly diminished, and given that patients who experience knee arthroplasty may even now experience impressive utilitarian disability postoperatively, the adequacy of physiotherapy after release is a legitimate inquiry. The present vulnerability with respect to viability makes it hard for charging associations, social insurance professionals, and patients to settle on choices in regards to such physiotherapy. We deliberately surveyed randomized controlled preliminaries to decide the adequacy of physiotherapy exercise after release regarding improving capacity, personal satisfaction, strolling, the scope of movement in the knee joint, and muscle quality for patients with

osteoarthritis after elective primary one-sided total knee arthroplasty.

Silman et al., (2001) stated that Osteoarthritis (OA) is unending degenerative disarray of multifactorial etiology depicted by loss of articular cartilage, hypertrophy of bone at the edges, subchondral sclerosis and extent of biochemical and morphological modifications of the synovial film and joint case. Obsessive changes in the late period of OA consolidate progressing, ulceration and focal separating of the articular cartilage; synovial exacerbation in like manner may occur. Ordinary clinical side effects are tormenting, particularly after deferred development and weight bearing; however, solidness is experienced after inertia.




Figure 1.2: Risk factors for osteoarthritis

Knee osteoarthritis (KOA) is a degenerative knee disease identified with pain, swelling, firmness, restricted ambulation, and declined parity work. It has been believed that chondrocytes experience awkward developing, which called "stress-initiated senescent express" which is the clarification behind cartilage degeneration. The provocative systems, the lessening of lubricin levels, and besides the impedances of the synovial fluid lubing up limit, which is immovably related to the headway of osteoarthritis have furthermore been acknowledged. It is a mileage joint inflammation result from the dreary pressure wounds of the joint and once in a while physical damage can worsen the circumstance; in any case, according to the Osteoarthritis Research International (OARSI) rules for the nonsurgical administration of KOA, exercise was recommended to improve the limit and exercises backing of people with KOA. [AAOS]

Patients with osteoarthritis (OA) of the knee are depicted basically by articular cartilage degeneration and an auxiliary peri-articular bone response. **Symmons D et al., (2000)** Around the globe, the regularity rate of OA is 9.6% for men and 18% for women >60 years. In India, OA is the second most ordinary rheumatologic issue and has a transcendence rate of 22 to 39. Clinically it presents as pain in and around the joint, joint firmness as a general rule after rest, crepitation and restricted joint advancements related to muscle deficiency. The most grounded risk factors for OA are age and hereditary qualities. Other danger components fuse female sexual orientation, rotundity, cigarette smoking, intra-articular breaks, chondrocalcinosis, valuable stones in joint fluid/cartilage, deferred immobilization, joint hypermobility, insecurity, fringe neuropathy, postponed word related or sports pressure.

Young-Joon Choi, and Ho Jong Ra (2016) proposed that Total Knee Arthroplasty (TKA) is a surgery used to mitigate pain and re-establish work in patients with serious disease of the knee joint, mostly knee osteoarthritis. As of late, there has been a significant improvement of Minimally Invasive Surgery procedures (MIS); these strategies for knee arthroplasty, presented with the point of decreasing

the harms to the extensor component, have developed quickly in the course of the most recent couple of years. MIS saves the tissues and the biomechanics of the knee joint, prompting better practical results with less postoperative pain, decreased blood misfortune with lower need of transfusions, an early rehabilitation with a quicker useful recuperation and a progressively fast accomplishing of utilitarian objectives.

There are confirmations of a connection between the postoperative planning of rehabilitation and the frequency of confusions like periprosthetic infections and profound vein thromboses, making rehabilitation a correlative technique to surgery. Results after TKA surgery should be identified with the power and kind of post-employable rehabilitation, with exposed proof of this relationship; in particular, rehabilitation protocols with early postoperative ambulation are exceedingly successful, prompting shorter postoperative hospital stays and lower pain relieving admission yet most examinations don't decisively depict rehabilitation protocols.

The achievement of the total knee strategy relies upon numerous elements, including quiet choice, prosthesis structure, the preoperative state of the joint, careful system (counting soft tissue adjusting and appendage arrangement) and postoperative rehabilitation.

Total knee replacement has progressed significantly since the main endeavours over a century back, when Theophilus Gluck planned and embedded a total knee made of ivory, balancing out the inserts with plaster of Paris and colophony (a translucent, fragile substance created from pine oleo sap, which is utilized in varnishes and inks). Gluck's total knee fizzled for an assortment of reasons, including poor bearing surface, inappropriate obsession, and continuous infections. With numerous enhancements, the Walldius pivot was presented in 1951. Made of acrylic, and later moved up to cobalt chromium (COCR) in 1958, this embed was utilized until the mid-1970s. In 1968, Frank H Gunston, MD, a Canadian specialist, planned the first polycentric knee. Two essential defects in his plan constrained its prosperity: it didn't supplant all the condylar surfaces, and it had a little contact zone. This polycentric knee was made of treated steel and just supplanted the weight-bearing piece of the knee, which is definitely not a genuine condylar knee. A restricted, polycentric metal supplanted the weight-bearing piece of the condyles, and the tibia was supplanted with tight, plastic sprinters. This took into account negligible revolution. Subsequently, these segments slackened sooner or later, which brought about the disappointment of this polycentric knee.

Risks Related to Total Knee Arthroplasty (Knee Replacement)

There are risks and inconveniences with this strategy. They incorporate yet are not constrained to the accompanying. General risks:

- Infection can happen, requiring antibiotics and further treatment.
- Bleeding could happen and may require an arrival to the working room. Bleeding is progressively normal in the event that you have been taking blood thinning drugs, for example, Warfarin, Aspirin, Clopidogrel (Plavix or Iscover) or Dipyridamole (Persantin or Asasantin).
- Small areas of the lung can collapse, expanding the danger of chest infection. This may require antibiotics and physiotherapy.
- Increased hazard in hefty individuals of wound infection, chest infection, heart and lung intricacies, and thrombosis.
- Heart attack or stroke could happen because of the strain on the heart.
- Blood clot in the leg (DVT) causing pain and swelling. In uncommon cases some portion of the coagulation may sever and go to the lungs.
- Death because of this system is conceivable.

2. REVIEW OF RELATED STUDIES

Hunter Warwick et al., (2019) contemplated that ongoing proof proposes an advantage to getting physical therapy (PT) a similar day as total joint arthroplasty (TJA), however generally little is thought about boundaries to giving PT in this obliged timeframe. We address the accompanying inquiries: (1) Are there demographic or perioperative variables related to getting deferred PT following TJA? (2) Does accepting quick PT following TJA influence transient results, for example, length of remain, release manner, or 30-day readmission? Methods. Primary TJA systems at a solitary focus were reflectively assessed. Quick PT was characterized as inside eight hours of surgery. Demographic and perioperative variables were analysed between patients who got prompt PT and the individuals who did not. We distinguished a properly coordinated control gathering of patients who got quick PT. Postoperative length of remain, release air, and 30-day readmissions were analysed between coordinated gatherings. Results. Altogether, 2051 primary TJA methods were explored. Of these, 226 (11.0%) got deferred PT. These patients had a higher rate of general anaesthesia (25.2% versus

17.8%, $p=0.006$), later usable begin time (13:26 11:31-14:38 versus 9:36 8:24-11:16, $p<0.001$), longer usable time (1.8 1.5-2.2 versus 1.6 1.4-1.8 hours, $p=0.002$), and higher by and large caseload upon the arrival of surgery (6 4-9 versus 5 4-8, $p=0.002$). A coordinated gathering of patients who got quick PT was distinguished. There were no distinctions in the postoperative length of remain or release attitude between coordinated quick and deferred PT gatherings, however postponed PT (OR 4.54; 95% CI 1.61-12.84; $p=0.004$) was related with a higher 30-day readmission rate. Conclusion. Boundaries to getting prompt PT following TJA included general anaesthesia, later usable begin time, longer usable time, and higher day by day caseload. These variables present potential focuses on improving the conveyance of prompt postoperative PT. Early PT may help decrease 30-day readmissions, however extra research is important to further describe this relationship.

Feng JE et al., (2018) Total knee arthroplasty (TKA) is the most usually performed inpatient surgery inside the USA and is evaluated to achieve 3.48 million techniques yearly by 2030. As worth based care activities keep on concentrating on hospital readmission rates and patient fulfilment, it has turned out to be fundamental for social insurance suppliers to create and execute a multidisciplinary approach to improve TKA results while minimizing pointless uses. Through this need, clinical care pathways have been created to standardize, sort out, and improve the quality and effectiveness of patient care while at the same time empowering the coordinated effort among different medical care suppliers. Here, we survey a few frameworks based projects and forte care rehearses that can be received into the standard orthopaedic practice.

Claudio Lisi et al., (2017) After TKA surgery should be identified with the force and kind of post-usable rehabilitation. Point of this paper is to depict our initial rehabilitation protocol following TKA with mini-obtrusive surgery in the prompt post-usable period and dissect useful recuperation and changes in pain scores in these patients. In this observational study, information was gathered on 215 total knee arthroplasty patients alluded to Orthopaedics and Traumatology inpatient ward from July 2012 to January 2014, treated with the equivalent solid start rehabilitation protocol. We recorded occasions to achieve utilitarian objectives (sitting, standing and helped ambulation) and pain after the treatment. Length of hospital remain in TKA was 4.6 ± 1.8 days, with a rehabilitation treatment length of 3.3 ± 1.3 days. The interim expected to accomplish the sitting position was 2.3 ± 0.7 days, to achieve the standing position was 2.6 ± 1.0 days to achieve the strolling practical objective was 2.9 ± 1.0 days. Pain NRS scores stayed beneath in the first and second post-employable day and underneath 3 from the third

post-usable day. End: Our study affirms that rehabilitation began when 24 hours after surgery with the mini-intrusive approach, empowers early virtualization of patients and early recuperation of strolling with decent control of pain.

Jeroen C et al., (2015) demonstrated that quick recuperation protocols diminish the length of hospital remain after Total Knee Arthroplasty (TKA) and Total Hip Arthroplasty (THA). Be that as it may, little is thought about the early postoperative stage. The motivation behind this study was to inspect which issues patients experienced during the initial a month and a half after primary TKA or THA surgery with quick recuperation. We welcomed twenty patients for a center gathering meeting that examined different subjects with respect to the initial a month and a half after hospital release. The center gathering gatherings were dissected subjectively. Patients were generally happy with the short length of hospital remain. Patients who lived alone needs more care and might want to remain longer in the hospital. After THA surgery all patients whined of failure to rest. More patients experienced pain after TKA surgery contrasted with THA surgery. All patients had different encounters with respect to physical therapy consequently a proof based rehabilitation protocol may be required.

Benazzo F et al., (2012) In particular, as of late, the mini-trivector approach was acquainted with broaden the advantages of the MIS strategies likewise in cases where it was hard to apply previously; indeed, one of the significant constraint for the utilization of MIS in knee arthroplasty is the poor versatility in case of real distortion or unbending nature of the knee joint and the mini-trivector approach for knee arthroplasties has been utilized as a legitimate option in contrast to great approaches, allowing to expand the advantages of MIS additionally to patients that wouldn't be reasonable for exemplary MIS procedures.

Bandholm T and Kehlet H (2012) Major surgery, including total hip arthroplasty (THA) and total knee arthroplasty (TKA), is trailed by a strengthening period, during which the loss of muscle quality and capacity is significant, particularly ahead of schedule after surgery. Lately, a mix of unimodal proof based perioperative care parts has been exhibited to upgrade recuperation, with diminished requirement for hospitalization, strengthening, and danger of medical confusions after significant surgery—the most optimized plan of attack system or improved recuperation programs. It is the idea of this procedure to deliberately and experimentally upgrade all perioperative care segments, with the general objective of improving recuperation. This is likewise the case for the care segment "physiotherapy exercise" after THA and TKA. The most recent meta-investigations on the effectiveness of physiotherapy exercise after THA and TKA by and a large reason that physiotherapy exercise after THA and TKA either does not work or isn't exceptionally

compelling. The explanation behind this might be that the "pill" of physiotherapy exercise normally offered after THA and TKA does not contain the correct dynamic fixings (too little power) or is offered at the off-base time (past the point of no return after surgery). We propose changing the concentration to prior started and increasingly escalated physiotherapy exercise after THA and TKA (quick track physiotherapy exercise), to decrease the early loss of muscle quality and capacity after surgery. In a perfect world, the physiotherapy exercise interventions after THA and TKA ought to be straightforward, utilizing few and well-picked exercises that are depicted in detail, holding fast to fundamental exercise physiology standards, if conceivable.

3. CASE STUDY

(A) The patient is a 53-year old female, who initially went to her primary care physician in 2004, at 46 years old, because of damage brought about by a fall at her workplace. She is a sprinter and dynamic. After examination, it is inferred that the patient does not have any mechanical side effects of locking. She has a negative Lachman test and a negative rotate move. She was sent for a MRI, which demonstrated a medial meniscal tear. X-beams show narrowing of the medial joint space, and plausible chondromalacia with imbuement. At the season of her underlying arrangement, the patient decided not to treat the tear, she was steady. Throughout the following five years, she was treated with various steroid shots, at that point painted with chlorhexidine gluconate arrangement and hung in the standard sterile design. The limit is exsanguinated and the tourniquet is swelled to 300 psi, and emptied before wound conclusion. Pins are put from horizontal to medial in the femur and the tibia for navigation. The midline entry point is made. Parapatellar arthrotomy is done, and in this case, a lot of clear joint liquid rose. The joint is uncovered and enrolled with Stryker 4.0. This patient has a thick, plain white synovial tissue all through the knee. It isn't in fronds and is increasingly membranous in nature. A synovectomy will be performed at the finish of the case.5 and a wipe, just as a disinfectant, because of pain in her knee. At that point, she was The Precision Knee Navigation System treated with a hyaluronate injection, which included a progression of three injections, two weeks separated. Along these lines, she encountered some alleviation. Following a half year, she rehashed the hyaluronate injection protocol. In June 2009, she selected a total knee supplanting with 4.0 navigation.

(B) This case related with rehabilitation of one 73-year-old male patient and intercessions used to enable him to come back to being practically autonomous. The patient had experienced right total knee arthroplasty and had recently

experienced respective total hip rehabilitation. Intercessions included manual therapy, physiotherapy, therapeutic exercise, interferential current and cryotherapy. The principle result estimates utilized for this situation were the quality, scope of movement, torment, and the Outpatient Physical Therapy Improvement in Motion Assessment Log Instrument. The patient was, in the long run, ready to make a full come back to his exercises in spite of not recovering the total scope of movement expected after total knee arthroplasty.

4. MATERIALS AND METHODS

Selection

We looked for randomized controlled trials of patients experiencing elective total knee arthroplasty for osteoarthritis who got an intervention of physiotherapy exercise after release from hospital. We utilized wide meanings of "physiotherapy" and "exercise" to incorporate any exercises or exercise program exhorted or given by physiotherapists or physical specialists during the rehabilitative period after release from hospital after surgery in the outpatient, network, or home setting. We avoided trials in which the intervention comprised of an electrical adjunct to physiotherapy, for example, the utilization of nonstop inactive movement. Physiotherapy exercise interventions included outpatient physiotherapy sessions and practical physiotherapy programs, in which exercises depend on utilitarian exercises. Trials were incorporated on the off chance that they examined a physiotherapy intervention contrasted and usual or standard care or looked at two changed kinds of applicable physiotherapy intervention. Usual or standard care alludes to the continuation of home exercise projects given to patients during a stay in hospital. These projects usually comprise of isometric or basic reinforcing exercises, exercises to recover the scope of development, and stretches. Effectiveness results were proportions of practical exercises of everyday living, strolling, self-revealed proportions of personal satisfaction, muscle quality, and scope of movement in the knee joint. As most trials utilize useful measures instead of explicit pain results, we did exclude pain as an effective result. Two commentators (CML and CS) surveyed and concurred on study qualification.

Validity assessment, data abstraction, and quality assessment

We created and steered an information extraction structure utilizing quality pointers from the CONSORT explanation and the CASP rules. Comparative investigation of individual quality parts has recently been utilized in surveys of physiotherapy and is supported to maintain a strategic distance from realized issues related to existing composite scores. Things could be set apart as truly, no, misty, or incomplete. Things were set

apart as yes just on the off chance that they completely and unequivocally met the point by point criteria spread out in the CONSORT standards. Two reviewers (CML and KB) autonomously separated the information. KB was concealed to the key subtleties of each paper and the degree to which veiling was effective was evaluated. The covering rates were 80% for creators, 20% for diaries, 80% for creator affiliations, and 80% for financing sources, all of which with the exception of the diary of distribution were viewed as effective. The level of understanding between reviewers was 69.09% (κ 0.524, intra class connection coefficient (2,1) 0.49, 95% confidence interval 0.30 to 0.63).

We settled introductory differences with respect to study quality by talking until an agreement was come to. The real difference was uncommon; usually, the contradiction was the more minor "yes" to "fractional/hazy" or "no" "too" "halfway/indistinct" and 100% understanding was gotten. A third commentator (CS) was accessible in case of agreement not being come to, yet this was not required. Where key study subtleties were missing or hazy we reached creators for additional data.

Quantitative information analysis

We did meta-analysis for knee work, strolling, the scope of joint movement, and personal satisfaction with R2.3.1 and the meta bundle. Our result was the score at the picked time point as opposed to the adjustment in the score as this amplified the number of equivalent investigations. The time focuses utilized were three to four months after surgery and a year after surgery. On the off chance that a similar measure was accounted for we utilized weighted mean contrasts, else we utilized standardized impact sizes (little (0.2), medium (0.5), and enormous (0.8)). We utilized fixed impact models and 95% confidence intervals all through and performed a trial of heterogeneity (χ^2) at a 5% significance level, however, we acknowledge these have low power since few investigations were accessible for meta-examinations. We are likewise determined to give an estimation of the level of heterogeneity between the trials in the meta-examination. Irregular impacts models were not considered as there was no convincing proof of heterogeneity and assessing the variation between studies is troublesome with such low numbers. The distinctions were determined so certain distinctions show that the impact favoured treatment and negative contrasts that the impact favoured control or usual care. We thought of it as wrong to evaluate distribution inclination as a result of the modest number of trials.

Table 1: Heterogeneity χ^2 and I2 test* results for range of outcomes

Outcome	Three to four months after surgery	12 months after surgery
Function	$\chi^2=0.76$, df=2, P=0.66	$\chi^2=2.37$, df=3, P=0.49
Walking	$\chi^2=0.52$, df=1, P=0.47	$\chi^2=0.40$, df=1, P=0.53
Joint range of motion	$\chi^2=1.45$, df=2, P=0.49	$\chi^2=2.31$, df=3, P=0.51
Quality of life	$\chi^2=0.39$, df=1, P=0.49	$\chi^2=0.89$, df=2, P=0.62

*I² =0% in all tests.

V. RESULTS

We distinguished and screened 27 possibly pertinent investigations. Of these, six investigations were incorporated into the efficient audit and five in the meta-analysis. Table 1 gives subtleties of prohibited examinations. Table 4 gives the consequences of the analysis of heterogeneity.

Measures of function (five trials)

Five of the investigations contained a proportion of function. The measures utilized incorporated the 12 thing Oxford knee score, which measures functional capacity, including pain, (scores 12-60, low score shows high function) the American Knee Society clinical rating score, which measures pain, development, steadiness, and functional movement (scores 0-100, high score demonstrates great); the 24 thing Western Ontario and McMaster Universities osteoarthritis record (WOMAC), which has areas for pain, solidness, and function (scored as a rate and out of 0-170 for function (low scores are positive); and the Bartlett patellar score, which measures anterior knee pain, quadriceps quality, and function (scores 3-30, high scores are ideal).

Inside the individual trials, three found no huge contrasts between gatherings. Ice et al found huge contrasts inside gatherings for the treatment arm, demonstrating an advantage of treatment. Some displayed no outcomes in their distributed conceptual however provided summary statistics for their results, allowing us to incorporate their study in the meta-analysis. A specialist found noteworthy contrasts between the two gatherings, for the intervention, at four and six months after arthroplasty however not at 12 months.

Figure 2 demonstrates the three investigations with information on functioning at three to four months and 12 months after surgery. Where studies included more than one proportion of function we chose to utilize the Oxford knee and the WOMAC scores as these incorporated all segment trials. No preliminary included both these scores. At three to four months the standardized impact size was 0.33 (95% confidence interval 0.07 to 0.58), which is viewed as small to moderate




Figure 1: Forest plot of standardized effect sizes with confidence intervals for function and results of test for heterogeneity

At 12 months, with one extra study, the impact size was near zero at -0.07 and the confidence interval (-0.28 to 0.14) included zero.

Walking (three trials)

Three knee arthroplasty trials utilized some type of a result measure for walking. The measures revealed included walking speed over a 10-meter separation, estimated in m/sec, and a six moment coordinated walking test, estimated in meters. The study by Moffet et al gave an account of time walking over a 50-meter walkway.




Figure 2: Forest plot of standardized effect sizes with confidence intervals for walking

The results from these trials were blended. One preliminary found no critical differences between groups, another discovered differences approaching significance, and the third preliminary found huge differences inside intervention groups. Figure 3 demonstrates that the intervention had no general impact on walking at either three or 12 months.

VI. CONCLUSION

After late primary total knee replacement, physiotherapy exercise interventions show short-term upgrades in physical function. This orderly survey offers help for the utilization of

physiotherapy exercise interventions with exercises dependent on functional exercises after release, as opposed to traditional home exercise and guidance programs, to acquire short term advantage after elective primary knee arthroplasty. There was a small to moderate standardized impact measure for functional exercise for function three to four months postoperatively. Small to moderate weighted mean differences, for functional exercise interventions, were seen for range of joint movement and personal satisfaction three to four months postoperatively. Any advantages seen after treatment did not endure to one-year follow-up.

REFERENCES: -

1. Silman AJ, Hochberg MC: 2nd ed. Oxford: Oxford University Press; 2001. *Epidemiology of the Rheumatic Diseases*.
2. Symmons D, Mathers C, Pflieger B (2002). Global Burden of Osteoarthritis in year 2000: Global burden of disease 2000 study. World health report. 2002;5 Version 2
3. Young-Joon Choi, and Ho Jong Ra (2016). "Patient Satisfaction after Total Knee Arthroplasty" *Knee Surg Relat Res* 2016 Mar; 28(1): 1-15 <https://doi.org/10.5792/ksrr.2016.28.1.1>
4. Hunter Warwick, Andrew George, Claire Howell, Cynthia Green, Thorsten M. Seyler, and William A. Jiranek (2019). "Immediate Physical Therapy following Total Joint Arthroplasty: Barriers and Impact on Short-Term Outcomes," *Advances in Orthopedics*, vol. 2019, Article ID 6051476, 7 pages, 2019. <https://doi.org/10.1155/2019/6051476>.
5. Feng JE, Novikov D, Anoushiravani AA, Schwarzkopf R (2018). "Total knee arthroplasty: improving outcomes with a multidisciplinary approach" *Journal of Multidisciplinary Healthcare* » Volume 11 25 January 2018 Volume 2018:11 Pages 63—73
6. Jeroen C. van Egmond, Hennie Verburg, Stephan B.W. Vehmeijer, Nina M.C. Mathijssen (2015) "Early follow-up after primary total knee and total hip arthroplasty with rapid recovery : Focus groups" *Acta Orthopædica Belgica*, Vol. 81 - 3 – 2015, pp-447-453
7. Ranawat C. SOA Textbook (2006). "History of Total Knee Replacement." The Southern Orthopaedic Association and Data Trace Publishing Company.
8. Benazzo F, Rossi SM (2012). The trivector approach for minimally invasive total knee arthroplasty: a technical note. *J Orthop Traumatol*; 13: pp. 159-62.
9. R. Development Core Team. R (2006). *A language and environment for statistical computing*. Vienna: R Foundation for Statistical Computing.
10. Impact of osteoarthritis: results of a nationwide survey of 10,000 patients consulting for OA. *Joint Bone Spine* 2005; 72: pp. 235-40
11. Noble PC, Gordon MJ, Weiss JM, Reddix RN, Condit MA, Mathis KB (2012). Does total knee replacement restore normal knee function? *Clin Orthops Rel Res* 2005;413: pp. 157-65
12. Bandholm T, Kehlet H (2012). Physiotherapy exercise after fast-track total hip and knee arthroplasty: time for reconsideration? *Arch Phys Med Rehabil* 2012; 93: pp. 1292-4.
14. Chan EY, Blyth FM, Nairn L, Fransen M (2013). Acute postoperative pain following hospital discharge after total knee arthroplasty. *Osteoarthritis Cartilage* 2013 ; 21: pp. 1257-63.
15. Minns Lowe CJ, Barker KL, Dewey M, Sackley CM (2007). Effectiveness of physiotherapy exercise after knee arthroplasty for osteoarthritis: systematic review and meta-analysis of randomised controlled trials. *BMJ* 2007; 335: pp. 812
16. Naylor JM, Harmer A, Fransen M, Crosbie J, Innes L (2006). Status of physiotherapy rehabilitation following knee replacement in Australia. *Physiother Res Int* 2006;11: pp. 35-47.
17. Van Peppen RPS, Kwakkel G, Wood-Dauphinee S, Hendriks HJM, Van der Wees PhJ, Dekker J (2004). The impact of physical therapy on functional outcomes after stroke: what's the evidence? *Clin Rehabil* 2004;18: pp. 833-62.
18. AAOS: American Academy of Orthopaedic Surgeons. *Treatment of osteoarthritis of the knee: Evidence-based guideline. 2nd edition*. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2013. May 18

Corresponding Author

Sharick Shamsi*

Senior Physiotherapist at Prince Sultan Military
Medical City, Riyadh KSA

sharickshamai@gmail.com