

# Heavy Metal Impact Assessment Study on Soil, Plants, Human Health on Microorganisms

Aditya S. Pawar<sup>1\*</sup> Dr. Krishan Pal Singh<sup>2</sup>

<sup>1</sup> Research Scholar, OPJS University, Churu, Rajasthan

<sup>2</sup> Associate Professor, OPJS University, Churu, Rajasthan

**Abstract – Heavy metals are among the most significant sorts of contaminant in the environment. A few strategies previously used to tidy up the environment from these sorts of contaminants, however a large portion of them are expensive and hard to get ideal outcomes. Since heavy metals in the environment have kept on expanding there is have to determine their levels in the environment for productive environment the executives, subsequently the need to determine the degrees of heavy metals in the soil in India.**

**Keywords: Heavy Metal, Soil Impact, Environment, Microorganism**

-----X-----

## 1. INTRODUCTION

Pollution is an overall issue and its potential in affecting soundness of the human populace is incredible (1). The effect of pollution in the region of stuffed urban communities and from mechanical effluents and vehicles has arrived at an upsetting size and is exciting open mindfulness (2). Extreme degrees of pollution are making a great deal of harm human and creature wellbeing, plants including tropical downpour timberlands just as the more extensive environment (1). Pollution is the reason for some sicknesses, which influence the old as well as the youthful and the enthusiastic and all creatures and plants (3). The WHO report calls attention to that twenty million kids overall experience the ill effects of pollution which has become basic in view of overpopulation (3,4). An expected 1.2 billion individuals drink unclean water which is the wellspring of water related ailments that slaughter between five-ten million individuals for the most part youngsters around the globe (5).

Heavy metals from a few sources have been noted to make huge pollution issues however investigation and evaluation are still lackillg to figure their effect on living life forms.

There are various kinds of pollution among which pollution brought about by lethal degree of heavy metal poisons is called heavy metal pollution (6). Heavy metals are components having a thickness more prominent than 5 g/cm<sup>3</sup> in their natural structure (6; 9). Human science is brimming with cases where heavy metal danger has prompted

mass passings (10). Every heavy metal are lethal to living beings at over the top fixations, yet some are fundamental for ordinary solid development and multiplication by plants at low yet basic focuses (6).

The heavy metals basic in follow components to plants incorporate Co, Cu, Fe, Mo and Zn and for creatures are Cr, Ni and Sn. The heavy metals Cd, Hg and Pb have not been demonstrated to be basic for either plants or creatures (9). It is imperative to note anyway that the groupings of individual metals in living tissues must be kept low and ought to be kept up inside thin breaking points to allow the ideal natural presentation of most life forms (9).

## 2. SOURCES OF HEAVY METAL POLLUTION IN THE ENVIRONMENT

Normally happening heavy metals are available in structures that are not promptly accessible for take-up by plants. They are commonly present in insoluble structures, as in mineral structures, or in hastened or complex structures that are not promptly accessible for plant take-up. Normally happening heavy metals have an incredible adsorption limit in soil and are hence not promptly accessible for living beings. These anthropogenic sources incorporate combination creation, environmental statement, battery generation, biosolids, covering, touchy assembling, inappropriate stacking of mechanical strong waste, calfskin tanning, mining, pesticides, phosphate manure, photographic materials, printing shades,

sewage water system, purifying, steel and electroplating enterprises, materials, and colors and wood conservation. Wellsprings of heavy metals, focuses in soil, soil properties, the degree and degree of take-up by plants, and the degree of ingestion by creatures, are the variables that impact the amassing of metal particles in the nourishment web.

### 3. MECHANISMS OF HEAVY METAL UPTAKE BY MICROORGANISMS

The cell structure of a microorganism can trap heavy metal particles and along these lines sorb them onto the coupling locales of the cell divider [11]. This procedure is called biosorption or aloof take-up, and is autonomous of the metabolic cycle. The measure of metal sorbed relies upon the active harmony and sythesis of the metal at the cell surface. The instrument includes a few procedures, including electrostatic association, particle trade, precipitation, the redox procedure, and surface complexation [12] (Figure1). The procedure is quick and can arrive at balance inside a couple of moments. Biosorption can be done by pieces of cells and tissues, or by dead biomass or living cells as inactive take-up by means of surface complexation onto the cell divider and other external layers [13].

#### 3.1 IMPACT ON SOIL

##### 3.1.1 Data analysis

Heavy metals impacts were not compelling up to soil, plants, and human prosperity yet what's more impact preparing the soil methodology by changing microbial tolerable assortment.

##### 3.1.2 Method of Soil Heavy Metal Pollution Assessment

The degree of soil HM pollution was overviewed as seeks after: first soil pH regards were characterized into three classes:  $<6.5$ ,  $6.5 \leq \text{pH} \leq 7.5$ , and  $>7.5$ ; second, the sullying edge for every earth HM was directed by means of land use (e.g., paddy fields) and pH class; third, the single defilement record (SPI) for each HM was settled (Equation (1)); finally, the Nemerow composite tainting list (NCPI) was determined (Equation (2)).

$$P_i = \frac{C_i}{S_i}$$

where  $C_i$  is the concentration of soil HM  $i$ , and  $S_i$  is the pollution threshold of  $i$ .

$$NCPI = \sqrt{\frac{(P_{imax})^2 + (\bar{P}_i)^2}{2}}$$

where  $P_{imax}$  is the greatest SPI estimation of every HM and  $\bar{P}_i$  is the mean SPI of every HM.

##### 3.1.3 Soil sampling

Sampling sites were chosen in line with anthropogenic sources of heavy metals. At each sampling point, approximately 0.5 kg of soil was collected 0-10 cm in depth using a stainless steel sampler.

##### 3.1.4 Sample digestion

Well mixed samples of 1 g each were gauged utilizing a scientech Zeta arrangement electronic equalization produced in the year 2000. The samples were placed into 250 ml glass measuring utensil and processed with 24 ml of water regia and afterward vanished to approach dryness. The soil samples were then disintegrated in 10 ml of 2% nitric corrosive, sifted and afterward weakened to 100 ml with refined water.

##### 3.1.6 Method Validation

The assimilation strategy and nuclear ingestion spectroscopy investigation were approved by recuperation technique. One gram of haphazardly chose soil powder was spiked with three distinct centralizations of heavy metals each in turn (1.0, 1.5, 2.0 ppm) each run in with the AAS 44 machine.

##### 3.1.7 Sample analysis

Buck logical (210 VGF) fire nuclear ingestion spectrophotometer machine was utilized in this investigation. Its parameters were set by the particulars given in the makes manual including light present and fuel arrangement of air/acetylene fire.

#### 3.2 Impact on plant

##### 3.2.2 Heavy metal determination:

The quantitative investigation of the heavy metals in plants was cultivated by instrumental methods, nuclear retention spectrometry (AAS), as indicated by principles global with a spectrometer Thermo Electron Model S Series AA SOLAAR, programming stage.

The heavy metal was determined from leaves utilizing the technique for Trivedi et al, (1987). One g dried plant material was taken into a 100ml Kjeldahl carafe, 1ml of 60% HC104, 5ml HN03 and 0.5ml H2S04 were included and warmed at low temperature. Warmth was expanded and material was processed for 10-15 minutes.

### 3.2.3 Study on soil

Toward the finish of the examination control and heavy metal tainted soils were dissected for EC, pH, Chloride, Carbonate, Bicarbonate, Calcium hardness, Magnesium hardness, Phosphorus, Potassium and Organic carbon.

### 3.2.6 Study on metabolism:

The third leaf (in repeat) of 15,30,45,60,75,90 and 105 days old control and treated plants were examined for starch digestion, protein digestion, antioxidative chemicals and IAA oxidase action, complete phenol.

## 4. IMPACT ON MICROORGANISM

The soil samples were collected in disinfected polyethylene compartments with the assistance of cleaned spatula. The samples for microbial examinations were taken at a profundity of 15 cm and moved to the research center for microbiological ponders. Three samples were taken for each duplicate and mixed equitably for metal and microbial investigation.

Modern sewage water samples: 25 ml of mechanical sewage water was taken in a tapered jar. It was processed by nitric: perchloric corrosive blend as portrayed in standard techniques (APHA, 1995).

### 4.1 Impact on soil

#### 4.1.1 Method validation

The systematic presentation of the strategy for investigation was finished utilizing recuperation test on certain samples to be examined. In every one of the counts of rate recuperation cases, the technique delivered a rate recuperation of between 95%-102% as appeared in Table 4.1 consequently making the strategy dependable for this examination.

#### 4.1.2 Heavy metals in agricultural and roadside soil

The Table 4.2 below shows the mean levels of the elements obtained in the four regions of Division 1 and division 2.


**Table 4.2: Heavy metals in different sub regions of two divisions of India in mg/kg**

Parameter	Town(n=60) Mean±SE	Village Areas(n=30) Mean± SE	crop growing area(n=60) Mean± SE	Livestock Area(n=24) Mean± SE	p-value
Cd	0.530±0.150	0.760±0.280	0.620±0.170	0.450±0.160	0.792
Cr	3.250±0.41b	1.460±0.270 a	3.480±0.480b	4.210±0.640b	0.008
Mn	19.480±1.220a	13.370±1.620 a	19.950±2.140a	26.830±3.290b	0.004
Pb	67.500±7.760b	31.000±6.930a	54.660±4.410ab	44.650±9.530 ab	0.005
Zn	403.300±44.340	322.440±108.990	305.990±83.030	165.990±16.82	0.0260

Mean ±SE values followed by same letters within the same row are not significantly different ( p-value 0.05, SNK test)

The mean degree of Cd in the soil inside the four chose sub locales, ran from 0.450-0.760 mg/kg. Town zone had the most elevated Cd mean of 0.760 mg/kg in the soil pursued by crop developing zone with a mean of 0.620 mg/kg. Division 2 domesticated animals region had the most reduced mean of 0.450±0.160 mg/kg. Cd displayed lower levels of defilement than those of different metals in this investigation. ANOVA showed no huge contrast between levels of Cd in soils from Town, villaage, crop developing zone and animals region (P=0.792).

An examination by Mico et al. (2006) on heavy metal substance of agrarian soils recorded comparative degrees of Cd in the soil going from 0.150-0.880 mg/kg. The prescribed scope of Cd in the soil is 0.070 – 1.100 mg/kg (Mico et al., 2006). Focus above 0.500 mg/kg mirrors the impact of human action (Mico et al., 2006). Figure 4.1 shows how the rate levels of Cd from the four sub districts inside the two divisions think about.



**Figure 4.1: Percentage Mean Cd levels in soils**

The mean degrees of Mn in the soil in the four sub areas extended from 13.370±1.620 to 26.830±3.290 mg/kg. Domesticated animals zone recorded the most noteworthy mean of 26.830±3.290 mg/kg and Town region recorded a mean of 13.370±1.620 mg/kg. town region and harvest developing region recorded comparable mean degrees of Mn in the soil. These degrees of Mn recorded in this durable were like those recorded by Al Yemen and Hashem (2006) in their investigation on heavy metals and microbial examination of soil samples. ANOVA at 95% certainty level uncovered a critical contrast between the mean degree of Mn in domesticated animals zone and the other three sub districts

(P=0.004). Fig 4.3 beneath analyzes the mean of Mn in soil in the four sub locales.




Figure 4.3: Mean Mn levels in soil from the four selected sub regions.

Soil for the most part contains 200 – 3000 mg/kg of Mn with a normal estimation of 600 mg/kg (Okunola et al., 2007). In the present examination, measure of Mn went from 13.37 – 26.83 mg/kg. The degrees of Mn in soils were generally low, suggesting mellow tainting of the metal in the soil. The mean degrees of lead around, town, crop developing zone and Livestock territory are 67.50±7.76, 31.00±6.93, 54.66±4.41 and 44.65±9.53mg/kg separately. Town had the most elevated mean of lead in the soil of the four sub areas. This could be credited to the surface spillover during blustery season why this conveyed to the dams and wells as 69 elevated levels are reflected in this water sources.

#### 4.2.1 Impact of heavy metal on growth and metabolism of plant (kalmegh)

Table 32: analysis of variance of enzymes and metabolites in the leaf of kalmegh grown without and with copper and cobalt as a soil pollutant

Factor	DF	COPPER			COBALT		
		MSS	F	CD (P=0.05)	MSS	F	CD (P=0.05)
INVERTASE ACTIVITY							
GP	6	9.36	4404.18**	0.02463	8.65	2894.95**	0.02921
T	4	3.14	1477.05**	0.02915	3.49	1169.50**	0.03456
R	1	0.004	1.68		0.04X10 <sup>-3</sup>	0.012	
GP*T	24	0.034	16.15**	0.01102	0.045	15.01**	0.01306
Error	34	0.002			0.003		
Total	69						
PROTEASE ACTIVITY							
GP	6	29.84	9287.96**	0.03029	29.29	11443.02**	0.02704
T	4	2.92	909.87**	0.03585	4.09	1598.28**	0.03199
R	1	0.004	1.24		0.002	0.805	
GP*T	24	0.037	11.51**	0.01355	0.042	16.35**	0.01210
Error	34	0.003			0.003		
Total	69						
PEROXIDASE ACTIVITY							
GP	6	0.035X10 <sup>-2</sup>	1428.94**	0.00027	0.035X10 <sup>-2</sup>	1771.84**	0.00024
T	4	0.034X10 <sup>-3</sup>	141.21**	0.00031	0.042X10 <sup>-3</sup>	212.70**	0.00028
R	1	0.018X10 <sup>-5</sup>	0.7481		0.057X10 <sup>-5</sup>	2.85	
GP*T	24	0.065X10 <sup>-6</sup>	2.60**	0.00012	0.049X10 <sup>-6</sup>	2.49**	0.00011
Error	34	0.025X10 <sup>-5</sup>			0.002X10 <sup>-6</sup>		
Total	69						
POLYPHENOL OXIDASE ACTIVITY							
GP	6	0.755X10 <sup>-4</sup>	28533.29**	0.000028	0.759X10 <sup>-4</sup>	31541.31**	0.000026
T	4	0.329X10 <sup>-5</sup>	1243.13**	0.000033	0.376X10 <sup>-5</sup>	1563.96**	0.000031
R	1	0.008X10 <sup>-5</sup>	3.12		0.002X10 <sup>-5</sup>	0.877	
GP*T	24	0.041X10 <sup>-5</sup>	15.59**	0.000012	0.037X10 <sup>-5</sup>	15.43**	0.000012
Error	34	0.003X10 <sup>-6</sup>			0.037X10 <sup>-7</sup>		
Total	69						

IAA OXIDASE ACTIVITY							
GP	6	9.49	4185.27**	0.02546	9.40	4377.40**	0.02477
T	4	2.83	1246.59**	0.03012	3.54	1648.06**	0.02931
R	1	0.01X10 <sup>-2</sup>	0.031		0.01X10 <sup>-3</sup>	0.06	
GP*T	24	0.084	36.95**	0.01139	0.052	24.23**	0.01107
Error	34	0.002			0.002		
Total	69						
NONREDUCING SUGAR CONTENT							
GP	6	257.18	413.50**	0.42155	224.66	1722.36**	0.19305
T	4	63.77	102.52**	0.49878	85.85	658.14**	0.22842
R	1	0.321	0.516		0.1142	0.8785	
GP*T	24	2.90	4.66*	0.18852	4.66	35.70**	0.08633
Error	34	0.622			0.13		
Total	69						


GP = GROWTH PERIOD

T=TREATMENT

\*\* = SIGNIFICANT AT 1%

\* = SIGNIFICANT AT 5%

Fig 32: IMPACT OF HEAVY METAL AS A SOIL POLLUTANT ON NONREDUCING SUGAR CONTENT OF KALMEGH



It appears that heavy metals smothered the sugar digestion. The heavy metal effect on parameters of starch digestion was in the request for decreasing sugar>non-lessening sugar>invertase movement. The huge diminishing in decreasing sugar level might be considered as biochemical side effects of heavy metals. Effect of heavy metal might be evaluated with the assistance of assurance of diminishing sugar in extremely youthful plant developed on heavy metal polluted soil.

### 5. IMPACT ON MICROORGANISM

It appears that heavy metals smothered the sugar digestion. The heavy metal effect on parameters of starch digestion was in the request for decreasing sugar>non-lessening sugar>invertase movement. The huge diminishing in decreasing sugar level might be considered as biochemical side effects of heavy metals. Effect of heavy metal might be evaluated with the assistance of assurance of diminishing sugar in extremely youthful plant developed on heavy metal polluted soil.

**Table 1 Concentrations of heavy metals in soil samples treated with Industrial sewage water quantitated by Atomic Absorption Spectrophotometer**

Soil Sample	Individual heavy metal content (mg/kg) expressed as mean value $\pm$ S.D.					
	Pb	Zn	Ni	Cr	Cu	Cd
SS1	427.5 $\pm$ 2.5	10812.5 $\pm$ 2.5	632.3 $\pm$ 2.1	184.5 $\pm$ 0.5	1467.5 $\pm$ 2.1	32.0 $\pm$ 2.0
SS2	180.0 $\pm$ 2.0	5212.3 $\pm$ 2.5	232.4 $\pm$ 2.6	70.1 $\pm$ 2.0	800.0 $\pm$ 2.5	13.5 $\pm$ 0.5
SS3	240.0 $\pm$ 2.0	6332.5 $\pm$ 2.5	570.0 $\pm$ 2.0	82.0 $\pm$ 2.5	1450.0 $\pm$ 2.5	20.7 $\pm$ 0.8
SS4	127.5 $\pm$ 2.5	3667.5 $\pm$ 2.5	200.0 $\pm$ 2.0	49.2 $\pm$ 2.7	140.5 $\pm$ 2.5	2.7 $\pm$ 0.2
SS5	68.5 $\pm$ 2.5	1667.5 $\pm$ 2.5	88.0 $\pm$ 2.0	38.5 $\pm$ 2.5	115.0 $\pm$ 2.5	1.2 $\pm$ 0.2
SS6	223.0 $\pm$ 2.0	5997.5 $\pm$ 2.5	400.0 $\pm$ 2.0	70.8 $\pm$ 2.5	1120.0 $\pm$ 1.5	14.7 $\pm$ 0.2
SS7	180.3 $\pm$ 2.5	4332.5 $\pm$ 2.5	150.0 $\pm$ 2.0	62.0 $\pm$ 2.0	632.0 $\pm$ 2.5	12.7 $\pm$ 0.2
SS8	107.5 $\pm$ 0.5	2917.5 $\pm$ 0.5	142.5 $\pm$ 1.2	38.5 $\pm$ 2.5	160.0 $\pm$ 2.5	1.5 $\pm$ 0.4
SS9 (control)	30.0 $\pm$ 2.0	96.0 $\pm$ 2.0	12.9 $\pm$ 2.5	15.0 $\pm$ 2.0	18.6 $\pm$ 2.1	N.D.

N.D: Not detected.

S.D: Standard deviation

**Table! Concentrations of heavy metals in industrial sewage water quantitated by Atomic Absorption Spectrophotometer**

Sample	Individual heavy metal content (mg/l) expressed as mean value $\pm$ S.D.					
	Pb	Zn	Ni	Cr	Cu	Cd
H1	109.5 $\pm$ 3.6	200.0 $\pm$ 2.0	140.5 $\pm$ 2.5	95.3 $\pm$ 2.5	110.0 $\pm$ 2.0	16.1 $\pm$ 2.0
H2	71.5 $\pm$ 2.5	74.3 $\pm$ 3.0	65.4 $\pm$ 4.5	68.0 $\pm$ 2.0	239.3 $\pm$ 2.5	6.6 $\pm$ 2.5

S.D: Standard deviation

## 6. CONCLUSION

The examination has demonstrated that there was impressive measure of heavy metals in ground and surface water; however the levels were underneath WHO most extreme reasonable levels for Mn, zinc and Cr. In hardly any cases the degrees of Cd surpassed the WHO greatest admissible point of confinement. The mean of lead was over the WHO most extreme passable points of confinement in ground and surface water from Division 1 and Division 2. Heavy metals invigorated peroxidase, polyphenol and IAA oxidase exercises. The phenol collection was higher in treated plants. Harmful, not all that basic and basic component caused pretty much comparative consequences for development and digestion of plants however poisonous component caused extreme impact even at low focus.

## 7. REFERENCES

1. K. Cho-Ruk, J. Kurukote, P. Supprung, and S. Vetayasuporn (2006). "Perennial plants in the phytoremediation of leadcontaminated soils," *Biotechnology*, vol. 5, no. 1, pp. 1–4.

2. Khan, A. and Ghouri, A. (2011). Environmental pollution: its effects on life and its remedies. *Journal of Arts, Science and Commerce*; 2: pp. 276-285.

3. Begum, A.; Ramaiah, M.; Harikrishna,; Khan, I. and Veena, K. (2009). Heavy metal pollution and chemical profile of Cauvery River water. *E-Journal of Chemistry*; 6: pp. 47-52.

4. Kanmony, C. (2009). Human rights and health care. New Delhi, India: Mittal Publication, pp. 73-76.

5. Pain, A. (2008). A brief look at lie: One man's view predicament. Leicester, UK. Troubador Publi limited pp. 81-84.

6. Bose, B. and Hemantaranjan, A. (2005). Developments in physiology, biochemistry and molecular biology of plants. New Delhi, India: New India Publishing Agency, pp. 105.

7. Voet, E.; Guinee, B. and Udode, H. (2008). Heavy metals: A problem solved?. Dordrecht, Netherlands: Kluwer Academic, pp. 4.

8. Shrivastav, R. (2001). Atmospheric heavy metal pollution, development of chronological records and geological monitoring. Accessed from [www.ias.ac.in](http://www.ias.ac.in).

9. Misra, S. and Mani, D. (2009). Soil pollution. New Delhi, India: S. B Nangia APH publishing corporation, pp. 29-59.

10. Yang, T.; Chen, M.-L.; Wang, J.H. (2015). Genetic and chemical modification of cells for selective separation and analysis of heavy metals of biological or environmental significance. *TrAC Trends Anal. Chem.*, 66, pp. 90–102. [CrossRef]

11. Fomina, M. & Gadd, G.M. (2014). Biosorption: Current perspectives on concept, definition and application *Bioresour. Technol*, 160, pp. 3–14. [CrossRef] [PubMed]

## Corresponding Author

**Aditya S. Pawar\***

Research Scholar, OPJS University, Churu, Rajasthan