

A Study of Some New Algebra Parameters Function and Their Applications

Anupama MPS*

Associate Professor, Department of Mathematics, Galgotias University, Greater Noida, Uttar Pradesh, India

Abstract – In a number of ways, from meteorological evaluations to monetary models, the exact display and expectation of the schedule progressively winds up. Besides the ability to anticipate an example is the need to establish an unequivocal quality or accuracy, preferably before the show opportunity occurs. This paper focuses on the question of predicting whether a number of reasonable models or much historical information are likely to present an opportunity. If all six models included in the information provided in this diagram are as accurate as they are, then anyone who analyzes the data is selected to describe what is likely to be the most extreme temperature every day, given six distinct expectations. For example, Medias, standard deviations and histograms are absolute sensitive choices to represent the likelihood of a temperature on a random day. If the information is severely skewed to the other side of the middle, this could lead to problems, in any case, with exceptions or too few information focuses to calculate a significant standard deviation or to create a valuable histogram.

Key Words – Algebra, Applications, Monetary Models, Information, Temperature Probability.

----- X -----

INTRODUCTION

This thesis entitled A Study on Recent Trends in Algebra is the zone of Modern Algebra which is one of the imperative and regularly developing parts of Mathematics. Algebra is personally identified with numerous parts of Mathematics, Engineering and different disciplines. In its quintessence can be depicted as the study of relations on delicate sets and On Precontinuous and Semi-preirresolute Functions, Preclosed Sets. The point of the present thesis is presenting sure new parameters in the Algebra and its ongoing applications and to study their Relationship with different various disciplines.

DEFINITIONS AND HIGHER TERMINOLOGY

Definition 1: Give X a chance to be a non-void Universal set. A fluffy subset A_n of X is a capacity $A: X \rightarrow [0,1]$.

Definition 2: Give G a chance to be a gathering. A fluffy subset A_n of G is said to be a fluffy subgroup of G on the off chance that it fulfilling the following maxims;

$$(i) A(xy) \geq \min \{ A(x), A(y) \}, \quad (ii) A(x^{-1}) \geq A(x) \text{ for all } x, y \in G.$$

Definition 3: Let A be fuzzy subset of a set X . For $t \in [0, 1]$, the level subset of A is the set,

$$A = \{ x \in X : \mu_A(x) \geq t \}.$$

This is known as a fluffy dimension subset of A .

Definition 4: Let A be a fuzzy of a group G . The sub group A of G . For $t \in [0,1]$ such that, $t \leq \mu_A(e)$ is called a level subgroup of A

Definition 5: A fluffy subgroup an of a gathering G is called fluffy ordinary if $A(x y) = A(y x)$

Definition 6: If $(G,)$ and $(G_1,)$ are two groups, then the function $f: G \rightarrow G_1$ is called a group homomorphism if, $f(x y) = f(x) f(y)$, for all x and $y \in G$.

Definition 7: Let X and X_1 be any two sets. Let $f: X \rightarrow X_1$ be any function and let A be a fuzzy subset in X , V be a fuzzy subset in $f(X) = X_1$, defined by $\mu_A(x)$, for all $x \in f^{-1}(y)$ $x \in X$ and $y \in X_1$. A is called a pre image of V under f and is denoted by $f^{-1}(V)$.

Definition 8: Let $f: G \rightarrow G_1$ be an anti auto morphism if $F(x y) = f(y) f(x)$, for all $x, y \in G$

Definition 9: Let μ is a fuzzy characteristic subgroup of a group G if $\mu f(x) = \mu(x)$

Note: Let $f: N \rightarrow N_1$ be a function and let μ and V be fuzzy sets in N and N_1 respectively. Then $f(\mu)$, the image of μ under f is under f is fuzzy set in N_1 defined by

$$f(\mu)(y) = \begin{cases} \sup \{ \mu(x) : x \in f^{-1}(y) \} & \text{if } f^{-1}(y) \neq \emptyset \\ 0 & \text{other wise} \end{cases}$$

For all $y \in N_1$. $f^{-1}(y)$ the pre image of y under f is a fuzzy set in N given by $f^{-1}(y)(x) = \mu(f(x))$ for all $x \in N$. Similarly to an α level cut, we have lower level cut as follows: Let μ be a fuzzy set in a set N . For $\alpha \in [0,1]$, the lower α level cut of μ is denoted $\alpha N_\mu = \{n \in N : \mu(n) \leq \alpha\}$.

Definition 10: Let (V,E) be an approximation space. Let X be a subspace of V . Then X is called primary if and only if

- (i) $\exists x \in V$ such that $X = \langle x \rangle$;
- (ii) $\forall y \in V$ if $X \subseteq \langle y \rangle$, then $X = \langle y \rangle$.

Definition 11: A subset A of a topological space (X, τ) is said to be preopen set if $\text{int}(\text{cl}(A)) \subset A$ or equivalently there exists $U \in \tau$ such that

$$A \subset U \subset \text{cl}(A).$$

The supplement of preopen set is called precloset set. The preclosure of $A \subset X$ is the intersection of all preclosed sets containing A and is denoted by $.p.\text{cl}(A)$

Definition 12: A space X is said to be connected if it is not the union of two non-empty disjoint open sets. Otherwise X is said to have a disconnection; i.e; X is disconnected, if there exist open sets A and B such that $A \neq \emptyset$, $B \neq \emptyset$, $A \cap B = \emptyset$ and $A \cup B = X$. Since A and B are also closed we may have a disconnection by closed sets.

Definition 13: A set Y is said to be preconnected if it is not the union of two non-empty disjoint preopen sets.

A subset $B \subset Y$ is said to be preconnected if it is preconnected as a subspace of Y . If $Y = A \cup B$ for some non-empty disjoint reopen sets A, B , then we say Y has predisconnection $\{A, B\}$.

Definition 14: A function: $f: X \rightarrow Y$ is said to be precontinuous if $f^{-1}(V) \in \mathcal{PO}(X)$, for every open set V of Y .

Definition 15: Let X be a non-empty Universal Set. A fuzzy Subset A of X is a function $A : X \rightarrow [0,1]$.

Definition 16: Let G be a group. A fuzzy subset A of G is said to be a fuzzy subgroup of G if it is satisfying the following axioms:

$$(i) A(xy) \geq \min \{A(x), A(y)\}, (ii) A(x^{-1}) \geq A(x) \text{ for all } x, y \in G.$$

Definition 17: Let A be a fuzzy subset of a set X . For $t \in [0,1]$ the level subset of A is the set, $A_t = \{x \in X : A(x) \geq t\}$. This is called a fuzzy level subset of A .

Definition 18: Let A be a fuzzy subgroup of a group G . The subgroup A_t of G , for $t \in [0,1]$ such that $A_t \neq \emptyset$ is called a level subgroup of A .

Definition 19: A fuzzy subgroup A of a group G is called fuzzy normal if $A(xy) = A(yx)$.

Definition 20: If (G, \cdot) and (G^1, \cdot) are any two groups, then the function $f : G \rightarrow G^1$ is called a group homomorphism if $f(xy) = f(x)f(y)$, for all x and $y \in G$.

Definition 21: We say that Ω is a germ of deformation of $\omega = f(X_1, \dots, X_n)$

$$dX_1 \wedge \dots \wedge dX_n \text{ if we have } F(X_1, \dots, X_n, 0) = f(X_1, \dots, X_n).$$

HISTORICAL REVIEW IN ALGEBRA

This deals with a survey of the summery literature and late application techniques in various united disciplines. Many understood theorems and results of world mathematicians, for example, Herstein I. N.35, Zuckerman H. S132, Grillet P. A.32, Kosoko B. what's more, Zadeh L. A.49 have been incorporated to render the work as independent as could be expected under the circumstances. Prior the Adelard of Bath and Naive concentrated the algebra interms of sets and groups. This exceptional background permitted the making of the concepts of ring and field. Other vital improvements were because of Sylow's, Jacobson, DeMorgans and Herstein I. N.35. Maybe a portion of the precursors utilized these concepts for advancing the science.

In the mid of the twentieth century numerous new concepts were produced, for example, Fuzzy set theory Zadeh L.A.127,128, harsh sets theory Pawlak Z.90 and their applications are numerous in Engineering Science and basic leadership prolems. Be that as it may, toward the finish of twentieth Century i.e. in 1999 Molodtsov D. A.81 introduced the concepts of delicate set theory as another arithmetic instrument for managing software engineering and technology. The first rot of the 21st century has opened another way to deal with this field of delicate Algebra. Numerous researchers over the world Are contributed a wonderful work on Algebraic structures of delicate sets and Singularities.


In the Ist mid rot of the 21st century: Szasz G. what's more, Griller P. A117, Fuchs I.27,28, Pandey S. K. what's more, Kare M.88,89, Warrack B. D. furthermore, Zadeh. L A125, Shabbir and Naaz113, Mefram Harpaul75 have enlivened the world with their gigantic work on singularities applications and delicate sets algebraic structure. Where as in the year 2009 delicate sets were utilized for investigation of evidence techniques in Discrete Mathematics, numerous regions of PC designing, data technology, physical sciences and their appropriateness in various Disciplines have opened another period.

The ordinary method for speaking to the solution to combinatorial issues was with an established set theory however as of late in the year 2012 and 2013, Samata B.95, Molodtsov D. A. what's more, Cagman A.80, Bosoki J.10, Shabbir and Cagman105 demonstrated an openness of delicate

sets is an option in contrast to established sets in algebra. These new hypothetical results and applications have roused us to work toward this path to contribute some new examinations. The vast majority of the genuine issues in sociologies, building, medicinal sciences, financial aspects and so forth the information included are loose in nature. The solutions of such issues include the utilization of numerical standards dependent on vulnerability and imprecision. Along these lines established set theory which depends on the fresh and correct case numerous not be completely reasonable for dealing with such issues of vulnerability. Also, in the year 2014 Shabbir and Naaz¹¹³ introduced delicate figurative mappings. Start of the year 2015 Mantu Saha and Baisnab A. B.⁶ are taking a shot at settled point theorem in delicate algebra with applications.

Algebraic geometry deals with peculiarity theory. Peculiarity theory is equivalent to Zeta Function, H Hyper Functions, Empirical process and Statistics.

- i) By utilizing this specific extension we can think about the conduct of any learning Machine dependent on the resolution of singularities.
- ii) The primary space of singularities is as appeared as follows



Further we have talked about the essential uses to date of the singularities lattice in the field of quantum science. The utilizations to which the frameworks have been put might be outlined into three primary classifications. In the first place, and most vital at present, has been the estimation of quantum-concoction parameters. We have considered in some detail the assurance of vitality eigenvalues, and a noteworthy piece of the work has been given to this assignment. In lesser detail we considered the assessment of a few different parameters, for example, bond orders. A few parameters have not been talked about by any stretch of the imagination, e.g. particle polarizabilities, despite the fact that these are promptly measurable from the parameters which have been considered. Further Abou-zaid S.¹ and Akram M.² have considered in various assortment of rings.

The mentioned parameters must be subordinate, rather than geometry, as an outcome b characteristic. In this way, even a parameter such as the bond length may be considered to be largely resolved by the species' peculiarity.

In setting quantum substance theorems, a second vital use of the peculiarity grid is. Although there was only a limited amount of work done here, the grid was used to create a number of theorems, such as the Pairing Theorem. It was used by Ruedenberg to create a number of theorems dependent upon a supposedly bonding arrangement as a parameter. Gutman and Trinajstić have shown that the Huckel rule is an explicit case of the wider Loop rule.

In studies which have shed light on the beginnings or parameters of quantum substances, the third use of the Network, which is currently largely unexplored. In this way, we can clarify how the Huckel system, with reference to the framework, unreasonably declines in own value. We showed how Huckel's theory is expanded into the 3-dimensional species by the framework of

peculiarity. In this way they illustrated the relationship between this framework and n-simplex characteristics. It should be evident that the use of the network to reveal an insight into the starting points of quantum science parameters and wonders can lead to numerous new and productive demands, in which sense it deserves more consideration than it has been up to now.

We have seen some arial surveys of the applications in celesterial mechanics with singularities on C^* algebra and in Julia sets. Further Arasu K. T.3 as yet watching some new results.

SEMI-PREIRRESOLUTE FUNCTIONS

Definition 1: A function $f : X \rightarrow Y$ is called semi-preirresolute if the inverse picture of every semi-revive set in Y is a semi-preopen set in X . Note that each semi-preirresolute delineate semi-precontinuous however not the opposite, which is appeared by the following example.

Example 1: Let $X = \{a, b, c, d\}$ and

$\tau = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{b, d\}, \{a, b, c\}, \{a, b, d\}, \{b, c, d\}, X\}; Y = \{m, n, l\}$
 and $\sigma = \{\emptyset, \{m\}, Y\}$. Let $f : X \rightarrow Y$ be a mapping defined by $f(a) = m$, $f(b) = f(c) = l$, and $f(d) = n$. Then, clearly f is semi-precontinuous but it is not a semi-preirresolute map since $f^{-1}(\{m, n\}) = \{a, d\}$ which is not a semi-preopen set in (Y, σ) .

We portray the semi-preirresolute mappings in the following theorem.

Theorem 1: The following statements are equivalent for a function $f: X \rightarrow Y$:

- (i) f is semi-preirresolute.
- (ii) For each point x of X and each semi-preneighborhood V of $f(x)$, there exists a semi-preneighborhood U of x such that $f(U) \sqsubseteq V$.
- (iii) For each $x \in X$ and each $V \sqsubseteq SPO(f(x))$, there exists $U \sqsubseteq SPO(x)$, such that $f(U) \sqsubseteq V$.

Proof: (i) \Leftrightarrow (ii). Assume $x \in X$ and V is a semi-preopen set containing $f(x)$. Then by (ii), there exists a semi-preopen set G such that $W = f^{-1}(V)$ be a semi-preopen set in X containing x and hence $f(W) \subset f(f^{-1}(V)) \subset V$.

(ii) \Leftrightarrow (iii). Assume that $V \subset Y$ is a semi-preopen set containing $f(x)$. Then by (ii), there exist is a semi-preopen set G such that $x \in G \subset f^{-1}(V)$. Therefore, $x \in f^{-1}(V) \subset cl(f^{-1}(V))$. This shows that $cl(f^{-1}(V))$ is a semi-preneighborhood of x .

(iii) \Leftrightarrow (i). Let V be a semi-preopen set in Y , then $cl(f^{-1}(V))$ is semi-preneighborhood of each $x \in f^{-1}(V)$. Thus, for each x is a semi-preinterior point of $cl(f^{-1}(V))$ which implies that $f^{-1}(V) \subset \text{int } cl(f^{-1}(V)) \subset cl(\text{int } cl(f^{-1}(V)))$. Therefore, $f^{-1}(V)$ is a semi-preopen set in X and hence f is a semipreirresolute map and we state the following theorems.

Theorem 2: If $f: X \rightarrow Y$ is a preopen and preirresolute mapping, then f is a semi-preirresolute. We note that every semi-preopen map is semi-preopen but not the converse further we state the following theorem.

Theorem 3: If $f: X \rightarrow Y$ is semi-preirresolute and $g: Y \rightarrow Z$ is semi-precontinuous, then $g \circ f$ is a semi-precontinuous map,

Theorem 4: If $f: X \rightarrow Y$ is continuous and open then $f^{-1}(clA)$ for every subset A of Y .

Theorem 5 : Let $f: X \rightarrow Y$ be a continuous open and preirresolute mapping, then f is a semi-preirresolute mapping.

Proof: Let $A \in SPO(Y)$, then there exists a preopen set $U \subset Y$ such that $U \subset A \subset clU$. Then by Theorem 17, $f^{-1}(clU) = cl(f^{-1}(U))$. Also, we have $f^{-1}(U) \subset f^{-1}(A) \subset f^{-1}(clU) = cl(f^{-1}(U))$.

Since f is a preirresolute map, then $f^{-1}(U)$ is a preopen set in X , and hence $f^{-1}(A)$ is a semi-preopen set in X . Thus, f is a semi-preirresolute map.

Theorem 6: A mapping $f: X \rightarrow Y$ is semi-preirresolute if and only if for every semi-preclosed set F of Y , $f^{-1}(F)$ is a semi-Preclosed set in X .

PRE-SEMIOPEN FUNCTIONS

We acquaint the presemipreopen mappings closely resembling with pre-semiopen mappings.

Definition 7: A function $f: X \rightarrow Y$ is called pre-semipreopen if the image of each semi-preopen set in X is semi-preopen set in Y .

We take note of that each semi-preopen delineate semi-preopen yet not the opposite.

Theorem: Let c and $g: Y \rightarrow Z$ be two maps such that $g \circ f$ is a pre-semipreopen map. Then,

- (i) If f is semi-preirresolute surjection, at that point g is a pre-semi-preopen outline.
- (ii) On the off chance that g is a semi-preirresolute infusion, f is a pre-semi-preopen delineate.

Proof : (i) let A be any semi-preopen set in Y , since f is a semi-preirresolute map, $f^{-1}(A)$ is a semi-preopen set in X . As $g \circ f$ is a pre-semi-preopen map and f is surjective, $g \circ f(f^{-1}(A)) = g(A)$, which is a semi-preopen set in Z . This implies that g is a pre-semi-preopen map.

- (ii) As we claimed in (i), we can prove the second part easily.

CONCLUSION:

In such cases, straightforward measurements won't give an extremely total image of the actual probabilities. This paper tends to such issues by asking: How would we be able to utilize devices, for example, principal segment analysis and free segment analysis to help precisely portray the probability of a future occasion dependent on a group of models for said occasion? To investigate this inquiry, we will test and develop a strategy proposed by Michael Dettinger, of the U.S. Geological Survey, for evaluating likelihood disseminations dependent on model

groups. This technique gives methods for assessing likelihood appropriations for time arrangement portrayed by small informational collections. At the point when there are just a small number of forecasts for a future occasion, there may not be sufficient information focuses to give an important image of the occasion's likelihood. Dettinger's strategy utilizes a principal part analysis (PCA) to partition the information into isolated segments, which are then redistributed so as to give an extensive number of new "forecasts" which pursue the patterns of the original however are presently sufficiently substantial in number to have the capacity to give progressively itemized data.

REFERENCES

1. Abou-zaid S (1991). On Fuzzy Subnear-ring and Ideals. Fuzzy Sets and Systems 44, pp. 139-146.
2. Akram M. and Dar K. (2007). On Anti Fuzzy Left h-ideals in Hemirings. Int. Math. Forum 2, pp. 2295-2304.
3. Arasu K. T. and Jungnickel (1990). Divisible difference sets with multiplier J. Algebra 33, pp. 35-62.
4. Arnold V. I. (1988). Singularities of smooth groups. Russ. Math. Sur 23. noj., pp. 1–42.
5. Arnold V. I. and Clent McCrory (1989). Singularities of smooth mapping. J. Russian Soviet. Math. 23, pp. 10-18.
6. Baisnab A. B. (2010). The radical and semi simplicity for arbitrary rings. Proc. Amer. J. Math. 17, pp. 300-320.
7. Balakrishnan R. and Suryanarayannan S. (2000). On P_k and P^k Near-rings. Bull. Malaysian Math. Soc. 23, pp. 9-24.
8. Bashir M. and Salim A. (2006). On soft sets and BCH algebra. J. Hac. Math. 39, pp. 205-217.
9. Biswas R. (1990). Fuzzy subgroups and anti-fuzzy subgroups. Fuzzy sets Systems 44, pp. 121-124.
10. Bosoki J. (2011). Classification of singularities in Linear Vector Spaces. Int. J. Comp. Sci. 12, pp. 439- 445.
11. Caell and Deye (1994). On the classification of discrete Fuzzy sets. J. Pattern recognition 26, pp. 461-470.
12. Chandrasekhara Rao K. and Swaminathan V. (2008). Anti-homomorphisms in Near-Rings. Inst. Jour. Comp. Sci. Math. 21 (2), pp. 83-88.

Corresponding Author

Anupama MPS*

Associate Professor, Department of Mathematics, Galgotias University, Greater Noida, Uttar Pradesh, India