

# The Concept of Relationship among Logic and Number Theory: Issues and Challenges

Vijaysinh Digambar Gaikwad\*

Assistant Professor, Dayanand Science College, Latur

## INTRODUCTION

In a prior period the applicable logical segment was recursion theory (decidability and undecidability). For  $\mathbb{Z}$ : the focal issue was Hilbert's tenth Problem, and the focal outcome is that recursively enumerable relations on  $\mathbb{Z}$ : are existentially quantifiable. The highpoint of determinability theory in  $\mathbb{Q}$  remains Julia Robinson's, that  $\mathbb{Z}$ : is  $n^3$ -perceptible in  $\mathbb{Q}$ . If  $\mathbb{Z}$ : is existentially determinable in  $\mathbb{Q}$  is obscure (in the event that it is, Hilbert's tenth Problem for  $\mathbb{Q}$  is undecidable).

Recursion theory is consequently exceptionally pertinent for the logic of worldwide fields and their rings of whole numbers. Conversely, demonstrate theory is substantially more applicable for the logic of neighborhood fields, and for those territories of number theory with a geometric perspective.

The by regional standards minimal consummations of number fields have all experienced fruitful model-theoretic breakdowns. Along these lines Tarski (1930's) gotten the established comes about on definitions in  $\mathbb{C}$  and  $\mathbb{R}$ , while not work the 1960's did Ax-Kochen-Ersov obtain comparable to comes about for  $p$ -adic fields (and for numerous Henselian fields). The consummations are all decidable, however these days one gives more criticalness to the perceptibility part of the above breakdowns. One acquires typical structures for definitions, connects between the geometry of the set and the type of its definition, and different uniformities for number of associated parts, and in  $\mathbb{Q}_p$  for the type of different integrals (cf. Loeser's course).

In the 1980's the ring of all arithmetical whole numbers was demonstrated (by means of a nearby to worldwide standard including prior work in mathematically shut fields with valuation) to have an extremely clear perceptibility theory, and specifically to have the simple of Hilbert's tenth Problem decidable.

In the 1960's and 70's there were a few improvements in unadulterated model theory that headed some an opportunity later to collaborations with number theory.

The leading was the work of Robinson school on model consummation, existentially shut structures, and compelling systems. The speculations of the fulfillments  $\mathbb{C}$ ,  $\mathbb{R}$ ,  $\mathbb{Q}_p$  all come up regularly in this settings. Be that as it may different hypotheses rise, without common models, yet which were to be enter segments in noteworthy connections in the 1990's. One is the theory of differentially shut fields (to be included in the Mordell-Lang guess, cf. the Pillay-Scanlon course), and an additional (not uncovered work 1990) is the theory Acfa of nonexclusive automorphisms (to be included in the "logical" approach to the Manin-Mumford guess). Different speculations of this sort identify with the lifting of Frobenius to the Witt vectors.

The other model-theoretic advancement, unquestionably deeper qua model theory, began with Morley's (1965) work launching model theoretic security. The 1965 paper gave a suggestive topological setting for first-request definability, and started a precise investigation of general ideas around the geometrical thoughts of measurement and autonomy. In spite of the fact that the following solidness theory applies just to  $\mathbb{C}$  around the culminations of number fields, recent day "neighborhood" forms of it have been included in latest cooperations of logic and number theory (cf. Pillay-Scanlon).

The other defining moment in the 60's was Ax's finalize the rudimentary theory of limited fields, where logic was seen to cooperate with Weil's Riemann Hypothesis for bends, and with Cebotarev's Theorem. The new speculations could be construed as fruitions of Robinson sort, and their determinability was suggestively dissected regarding Galois Stratification (cf. Loeser's addresses). In addition, one was soon prompted model theoretic inquiries regarding total Galois bunches, and those are identified with a dream of Grothendieck (see Pop's addresses). (This is in no way, shape or form the main situation where plans of Grothendieck, the "logician" of Bourbaki, have had crucial logical substance.)

As of recently in Ax-Kochen-Ersov one had seen the force of the drive "Let  $p$  head off to 0." After Ax's work one had wealthier situations for this thought.

Specifically, one could steadily approach the investigation of the Frobenius  $x^p$  as  $p$  has a tendency to 0. Here one reaches the Weil Conjectures and the chorological routines utilized within their evidence. Old subjects of Robinson, on limits in the theory of goals in polynomial rings, return in the more extensive setting of Intersection Theory and Weil Cohomology, and identify with the Grothendieck Standard Conjectures.

A different excellent guess that of Schanuel on transcendence of qualities of the intricate exponential, has as of late started to interface with logic. It was first seen in association with the decidability of the true and  $p$ -adic exponentials, and all the more as of late in a significant determinability theoretic study by Zilber of the undecidable complex exponential. Zilber's work interfaces regularly with diophantine geometry, and with old work of Ax (cf. Pillay-Scanlon).

### THEORY OF FINITE FIELDS

The decidability of the primary theory of limited fields was a noticeable problem around 1960, and the decidability of the relating issue for  $p$ -adics was verifiably seen to decrease to this by the Ake examination. Note that this was after the negative result of Hilbert's tenth Problem. One obviously knew counterexamples to local/global standards for diophantine mathematical statements, and may subsequently suspect a limitless distinction between the all inclusive theory of the  $p$ -adics and the general theory of the whole numbers. In 1963 Nerode demonstrated the decidability of the all inclusive theory of the  $p$ -adics by utilizing the reduction of the ring of  $p$ -adics whole numbers. His evidence was for a solitary  $p$ -adic, and no generalization to all  $p$ -adics at the same time was evident.

Hatchet acknowledged the general ultraproduct of limited fields, with a perspective to understanding the essentially all theory of limited fields. His generally striking perception was that Weil's profound result, the Riemann Hypothesis for bends over limited fields (or, rather, the resulting Lang-Weil evaluations) might give the prevailing adage for the nonprincipal ultraproducts. Ersov had located some unique instances of this, as well. The resulting Weil Axiom Scheme, together with a Galois-theoretic plan, totally axiomatized the ultraproducts. While the Galois adage is accurate for all limited fields, the Weil Axiom is accurate for none, however materializes in the breaking points furnished by ultraproducts.

Uniformity crosswise over Finite Fields : Ax's axiomatization relies on upon a nontrivial uniformity:

There exists a capacity  $F$  from  $N$  to  $N$ , whose careful shape is superfluous, in any event in primary scenarios, such that if  $V$  is a totally irreducible relative bend over  $F_g$  and  $g \geq F$  (genus( $v$ )) then  $V$  has a  $\mathbb{F}_q$  esteemed focus.

Presently, the vital focus is that the variety of a bend is limited above by a basic capacity of the level of polynomials demarcating the bend, freely of the coefficients of these polynomials, and free of the encompassing field. All the more usually and uniquely, if  $V$  has a place with a group of mixed bags (recorded, say, by a constructible set) there is a capacity  $G$  of the family with the goal that if  $g \geq G((V))$  then  $V$  has a  $\mathbb{F}_q$  esteemed focus. It accompanies effectively by the Los Theorem that the nonprincipal ultraproducts fulfill the property now reputed to be Pac, or (in my idea better) normally shut. This is :

(Pac) Every completely irreducible assortment has a focus.

This was honed by Geyer, who saw that it is sufficient to request that each completely irreducible bend has a focus.

Aphorisms : To make the above as a primary axiomatization, one should realize that the property completely irreducible is first request, that is has a definition not depending on coefficients or encompassing field. there are numerous approaches to see this, equally exceptional unless one has productive slants. For instance, one can utilize the effect, from the Robinsonian theory of limits in polynomial standards, that prime is primary, joined together with Tarski's quantifier-disposal for logarithmically shut fields.

To finish the axiomatization for the nonprincipal ultraproducts, one imposes two different conditions. One is the clear one, that the fields are impeccable, as limited fields may be. The other maxim plan is more huge. We know, by numbering, and utilizing the Frobenius automorphism of limited fields, that every limited field has precisely one development of every limited size. Checking has no clear convenient rudimentary variant, yet one can utilize antiquated formed variable based math (Tchirnhausen changes and the like) to show that the property of a field to have exactly one augmentation of every measurement (called quasifinite by Serre) is rudimentary.

Putting Pac, immaculate, and quasifinite together one gets a set of maxims for what we now call pseudofinite fields. There is by now a rich theory of these structures. One may have had reservations about the starting points of the aforementioned fields, in the farout universe of ultraproducts (Mumford) , yet their manifestation in such a variety of imperative comes about throughout the most recent 40 years has without a doubt secured their qualifications. They have showed up "in nature" since, in some ways. Jarden demonstrated that a non specific component of the Galois gathering of a countable Hilbertian field has settled field pseudofinite. Much later van nook Dries watched that the altered fields of an existentially shut contrast field is pseudofinite. Later still, Pop

demonstrated that the field of completely legitimate logarithmic numbers is Pac.

Galois Aspects : What I like about customarily shut is that it uncovers a model theory (of fields) for the class of general embeddings (which incorporates the classification of primary maps).the consistently shut fields are just the (Robinsonian) existentially shut structures for the classification of normal maps.

The model theory of standard maps is quite rich, on the grounds that it has a double model theory of Galois aggregations. regular maps of fields instigate (contravariantly) epimorphisms of supreme Galois bunches. The model theory of profinite assemblies

is a completely a significant number sorted undertaking, with no quantification over aggregation components, yet rather an entire run of "limited " quantifications over limited remainders. Its utility depends substantially on Herbrand's backwards limit development. Limitedly produced profinite bunches assume the part of limited assemblies in the comodel theory, and the Iwasawa or Frobenius amasses act like homogeneous models.

The Galois assembly of a customarily shut field is liable to a major cohomological constraint, specifically that it is projective. The abundance of the theory relies on upon the way that this idea is first-request (for the field), and has numerous comparable details. The pseudofinite fields have Galois bunch "z" and this gathering is limitedly created.

Hatchet indicated that each pseudofinite field is rudimentarily identical to a ul-traproduct of limited fields. For this, in trademark zero, Cebotarev's Theorem is utilized.

Hatchet utilizes the tried and true isomorphism approach to rudimentary comparability, and his verification yields that two pseudofinite fields are rudimentarily proportional if and just in the event that they have the same trademark and the same "supreme numbers", i.e the same monic polynomials, in one variable over  $\mathbb{R}$  , are reasonable in each. It is suggestive to give this a more invariant, less syntactic, detailing. Until one fixes a logarithmic conclusion of the prime field, the idea of logarithmic numbers has small sense. Actually, what Ax is appending to a theory of pseudofinite fields, say in trademark zero, is a conjugacy class of shut procyclic subgroups in  $\text{Gal}(q)$ . Besides, his investigation shows that this duty outlines a homeomorphism structure the Tarski space of complete hypotheses of pseudofinite fields of trademark zero to the Vietoris space of conjugacy classes of shut procyclic subgroups of the minimized gathering  $\text{Gal}(q)$ . James Gray has explained this extensively to fit the whole Tarski space, with no

confinement on trademark, into a Vietoris space joined to  $\text{Gal}(q)$ .

From these contemplations (however with less reflection than utilized above) Ax promptly demonstrates decidability of the theory of pseudofinite fields, and afterward, by thoughtfulness regarding the type of his maxims, decidability of the theory of limited fields.

## **BACKGROUND OF TARSKI, MAL'CEV AND ROBINSON**

Tarski -Tarski, in the 1930's, committed:

- 1) Set-theoretic establishments of model theory, permitting exact meanings of structures, their grammar and semantics (not whatsoever bound to first-request semantics);
- 2) The establishments of perceptibility theory in the requested field  $\mathbb{R}$  ;
- 3) (Via Presburger) the establishments of perceptibility theory in the requested assembly  $\mathbb{Z}$ .

Later, in Berkeley, Tarski (and people) separated the essential morphisms (elementary embeddings), the Limit Theorem, and the ultra product development (and more general items, numerous significant to number theory). Szmielew made a major commitment by a deliberate examination of the first-request theory of abelian gatherings (however not yet going the extent that a disposal theory). But here, as in the majority of the following work of the Tarski school, the attention turned to decidability. In [40] a efficient examination is made of undecidability, for both finish and deficient theories. The system for vital undecidability is unmistakable here. There followed, a decade later, the still functional upgrade from the Mal'cev school. Forty years after the fact, very nearly all the open issues said there have been replied, or, all the more vitally, have been sent out to the universe of perceptibility. Lamentably, however, there is confirmation that huge numbers of the significant elucidations given there are new to the more youthful eras.

Most likely the deepest work done in Berkeley on logic and number theory was that of Julia and Raphael Robinson. The previous gave a  $\mathbb{Z}$  meaning of  $\mathbb{Z}$  in the ring  $\mathbb{Q}$  (never enhanced), and roused the research that built up and finally finished in 1970 in Matejasevic's negative result of Hilbert's tenth Problem for  $\mathbb{Z}$ . It is vital that Julia Robinson utilized local/global contemplations within her work, and that in (generally variants of) Matejasevic's evidence one uses the standard manifestations of quadratic enlargements of  $\mathbb{Q}$  .

Poonen's addresses are obviously committed to Hilbert's tenth Problem, and the fundamental issue of the undifferentiated from consequence for  $\mathbb{Q}$  and other worldwide structures.

Mal'cev: In 1936 he gave the strategy for graphs, and the general completeness/compactness hypothesis, and recently gave some quite creative provisions to gathering theory.

The school he established at Novosibirsk generated Ersov and Zilber around numerous others. these two are singled over here due to their remarkable contributions to the subject of our gathering (Ersov on p-adic and normally shut fields, bury alia, and Zilber on geometric model theory and diophantine geometry).

Abraham Robinson: He strove to open ways (both routes) between logic and polynomial math (thus number theory). For an extremely adept remark on this picture the stress on model-finish speculations, and Robinson's Test, in the shape very nearly of a Nullstellensatz, launched an advancement that has endured fifty years (and prospers still, as a result of an advantageous interaction with the geometric model theory began by Morley in a Tarskian setting).

Around Robinson's accomplishments, by these methods are:

- 4) Very applied approach to true shut fields and Hilbert's seventeenth Problem on wholes of squares(with limits);
- 5) Definability theory for logarithmically shut fields with valuation, critical thirty years after the fact in a local/global setting;
- 6) Bounds in polynomial goals, a theme as of now advancing in view of the requirements of logic of cohomology;
- 7) Factorial compactification in nonstandard dissection;
- 8) Differentially shut fields, an actually regular theory with no truly natural models (its ensuing worth is that it furnishes a rich "geometrical" finishing for diophantine geometry over capacity fields);
- 9) Generic structures by a mixed bag of "compelling" strategies, and the attention on finding sayings for those structures.

**SOAKED MODELS AND ULTRAPRODUCTS**

These originated from the Tarski school, and have demonstrated, notwithstanding their set-theoretic trappings, exceptionally helpful in connected situations,as a setting for changing over quantifier-end

comes about into outcomes on augmentation of isomorphisms. In the early days the technique was bound up with the ultra product construction, both in Keisler's work and in Kochen's paper which remains an excellent introduction to model-theoretic variable based math.

**P-ADIC FIELDS**

After 1964, one had a web of analogies interfacing the logics of the fruitions of number fields. I give a revisionist record. I was starting research in this period, and was impacted by numerous drives, for example Morley. Forty years on, there does general understand that Morley's plans have a place with those of the Tarski-Robinson-Mal'cev custom, however, actually, they don't make a difference straight to the crux speculations.

Before 1964 one comprehended, at first by means of Tarski's barehand strategies, the essential metamathematics of logarithmically shut and true shut fields, and no others. A mixed bag of techniques had , since Tarski, been conveyed on the essential speculations for instance, those of Robinson, Kochen, and the Shoenfield standard utilizing soaked models). I propose that one recollect that all these plausible outcomes.

Starting now, the fundamental characteristic is an uniform determinability theory (dependent upon an uniform quantifier-elimination)for the nearby fields originating from traditional number theory. Those fields are all generally reduced (see Weil's book for the unifying nature of this thought alone), and, in all cases however that of  $\mathbb{C}$  (a base for) the topology is arithmetically definable.so it is characteristic, first time adjust, to take a gander at these fields in the immaculate dialect of field theory.

Defining the topology - In the real field, the order (and hence a basis for the topology) is definable thus:

$$x \geq y \iff x - y \text{ is a square}$$

In a finite extension of  $\mathbb{Q}_p$ , the valuation ring (and hence the topology, is definable thus:

$$v(x) \geq 0 \iff 1 + \pi x^2 \text{ is a square}$$

where  $\pi$  is a uniformizing element, i.e an element whose value is minimal positive in  $\text{val}(K)$ . (Note that the use of  $pi$  can be eliminated by a standard trick of quantifying over possible uniformizing parameters).

When  $p \neq 2$ , one has to modify the definition thus:

$$v(x) \geq 0 \iff 1 + \pi x^8 \text{ is a square}$$

Now is a good moment to introduce the power predicates  $P_n$ , with the defining condition:

$$P_n(x) \iff x \text{ is an } n\text{-th power.}$$

So we have shown, uniformly for the real and the  $p$ -adic cases, that the topology is quantifier-free definable in terms of the  $P_2$ . The other power predicates are needed in the  $p$ -adic cases for quantifier-elimination. Note their link to Presburger arithmetic, since the valuation of an  $n$ -th power is divisible by  $n$ . The quantifier-eliminations in the  $p$ -adic fields somehow reflect (among other features) the quantifier-elimination in the value group.

It is often more natural to use modifications  $P_n^*$  interpreted as the set of nonzero  $n$ -th powers (for the same reason as it is more natural to use strict less than rather than less than or equal in the real case).

That the topology of  $\mathbb{C}$  is not field-theoretically definable comes from the existence of discontinuous field automorphisms of the complex field (the sort of phenomenon that distresses Deligne in .

An important analogy between the three families of fields  $\mathbb{K}$  (complex, real and  $p$ -adic) is that  $\text{Gal}(K)$  is prosolvable, and topologically finitely generated. Moreover, every finite extension of  $K$  is generated by an element of  $\mathbb{Q}^{alg}$ .

The fundamental result connecting the three definability theories, most illuminating, at least on first reading, if restricted to the cases of the complexes, the reals and the unramified developments of  $\mathbb{Q}_p$  (the ramified case is a spot harder,) is that each determinable connection is in the Boolean polynomial math created by the arithmetical sets and the sets demarcated by conditions

$$P_n^*(f(\bar{x}))$$

The primary cause behind my utilization of these predicates in the  $p$ -adic case was that taking forces has a considerable measure to do with building feasible augmentations. For  $\mathbb{C}$  there are no arithmetical amplifications, and all the force predicates are redundant for  $\mathbb{R}$  there is just a cyclic growth of request 2, concentrating a square foundation of  $-1$ . For the  $p$ -adics, all power predicates are required.

An amusing, but not accidental, observation is that the unit ball in the reals is perceptible by  $P_{2(1-x^2)}$ .

This shows an uniformity in definition with those for the  $p$ -adic unit balls, now taking  $p = -1$ . In fact, it prescribes interpreting the reals as the  $-1$ -adic numbers.

## CONCLUSION

It is educational to perceive how the state of the center aphorism frameworks has modified in excess of 40 years. Robinson's adages were dependably figured as far as illuminating comparisons in a solitary variable, and this sort of detailing won through the Ake revolution. For consistently shut fields, one was obliged to utilize adages about higher-dimensional mixtures, however Geyer permitted one to confine to planar bends. At different times in the 70's I taken a gander at growths of the theory of differentially shut fields to different hypotheses of fields with inference, and I discovered it regular to define relating aphorisms regarding mixed bags of higher extent and their "changes" under the determination. A more sumptuous form of this was displayed by Pierce and Pillay in.

The sayings for Acfa are of this shape too, giving consistency conditions for the multidimensional chart of the automorphism sigma to meet a subvariety of the result of a mixed bag and its "convert". Furthermore Zilber's maxims are of this shape too, for exp, yet with a gently infinitary flavour.

There are different situations where one has given nitty gritty met mathematical analyses without utilizing such formulations, but where it might well be important to continue to look for such adages. Clear illustrations are the true exponential field, and the Witt Frobenius.

## REFERENCES

- Baldwin J. T. and Lachlan A. H. (1971). On strongly minimal sets. *J. Symbolic Logic* 36, pp. 79–96.
- Belair L., Macintyre, A.J. and Scanlon, T. (1980). Model theory of Frobenius on Witt vectors.
- Chatzidakis, Zo'e ; van den Dries, Lou ; Macintyre, Angus . Definable sets over finite fields. *J. Reine Angew. Math.* 427 (1992), pp. 107–135.
- Denef J. (1988). van den Dries, L.  $p$ -adic and real subanalytic sets. *Ann. of Math.* (2) 128, no. 1, pp. 79–138.
- Hrushovski Ehud (2001). The Manin-Mumford conjecture and the model theory of difference fields. *Ann. Pure Appl. Logic* 112, no. 1, pp. 43–115.
- Lang S. (1970). *Algebraic Number Theory*, Addison-Wesley, New York.
- Macintyre (1986). Angus Twenty years of  $p$ -adic model theory. *Logic colloquium '84* (Manchester,

1984), 121–153, Stud. Logic Found. Math., 120, North-Holland, Amsterdam.

Macintyre Angus and Wilkie A. J. (1996). On the decidability of the real exponential field. *Kreiseliana*, 441–467, A K Peters, Wellesley, MA.

Marker (2002). *David Model Theory: An Introduction*. Graduate Texts in Mathematics 217, Springer Verlag.

Pop Florian (2002). Elementary equivalence versus isomorphism. *Invent. Math.* 150, no. 2, pp. 385–408.

Robinson Julia (1963). The decision problem for fields. 1965 *Theory of Models (Proc. Internat. Sympos. Berkeley)* pp. 299–311 North-Holland, Amsterdam

Skolem Th. (1955). Peano's axioms and models of arithmetic. *Mathematical interpretation of formal systems*, pp. 1–14. North-Holland Publishing Co., Amsterdam.

Tarski Alfred (1948). *A Decision Method for Elementary Algebra and Geometry*. RAND Corporation, Santa Monica, Calif., iii+60 pp.

Wilkie A. J. (1989). On the theory of the real exponential field. *Illinois J. Math.* 33, no. 3, pp. 384–408.

---

### Corresponding Author

**Vijaysinh Digambar Gaikwad\***

Assistant Professor, Dayanand Science College, Latur

E-Mail – [saritakhadse03@gmail.com](mailto:saritakhadse03@gmail.com)