



*Journal of Advances and
Scholarly Researches in
Allied Education*

*Vol. V, Issue X, April-2013,
ISSN 2230-7540*

**STUDY ON WATER SHADE PROJECT AND ITS
IMPORTANCE IN HARYANA**

Study on Water Shade Project and Its Importance in Haryana

Dr. Randhir Singh

Assistant Professor Govt. College, Bhiwani

Abstract – In spite of the favorable conditions to grow horticultural crops throughout the year in Haryana production is restricted because of water deficit. This is particularly noticeable during the dry season (December to June) when evaporation is higher than rainfall. Heavy rains in Haryana are concentrated in the wet season (July to August) and the average rainfall in St Croix is 1041 mm per year which corresponds to 60% of the pan evaporation. This water deficit can be ameliorated in the horticulture industry by growing crops under shady conditions. Plants require substantially less water (40% to 60%) because of reduced evapo-transpiration under shade conditions. In addition, many tropical high value ornamentals perform better in shady conditions. This project proposes to develop a water budget for shade-house production of ornamentals. The specific objectives of this study are: a) to determine the optimal WP of anthuriums by quantifying the shading effect on crop evapo-transpiration (ETc), growth and flower production; b) to determine the crop coefficients that associates ETc under shade with pan evaporation and potential evapo-transpiration (ETo); c) to estimate irrigation water requirement throughout the year for anthurium cut flower production under optimal shade conditions. A greenhouse with three shading levels (treatments) will be used for this study. Potted anthurium, a tropical ornamental cut flower, will be grown under each treatment and monitored for soil moisture and water used. ETc, plant growth and cut flower yield will be evaluated to determine the shading condition for optimal water productivity and water requirements. Climatic data (rain, relative humidity, pan evaporation, light intensity, temperature, etc) from a weather station will be analyzed to determine ETo. Water requirements under optimal growing conditions will be associated with ETo and pan evaporation to determine crop coefficients. The information obtained in this project will help horticulturists Haryana develop decision criteria for crop selection and adoption of a water-sustainable production system.

Key words: horticultural crops, Irrigation, potential evapo-transpiration, optimal growing conditions.

----- X -----

INTRODUCTION

Watersheds need a helping hand—yours

We all live in a watershed. At home, at work or at your child's school, chances are that water is trickling and gushing underfoot, flowing as waste.

Our [watersheds](#) connect us to each other and to the entire Bay Area. Everything we do—from mountaintop to marsh—affects our water and wildlife, our air and soil quality, our families and communities.

But our ties to local water resources have unraveled. Birds and fish have fled in the face of urban and industrial development. Shorelines sprout trash instead of grasses.

Join The Watershed Project in restoring the natural wonder of Haryana—one creek, marsh and watershed at a time. A bay simply holds water, an estuary mixes it. It's the difference between a glass of neat gin and a dry martini—gin is well and good but dry martinis are

holy, and though bays are useful, estuaries are explosive creators of life.

When one tugs at a single thing in nature, he finds it is attached to the rest of the world." John Muir

PROBLEM AND RESEARCH OBJECTIVES

In spite of the favorable conditions to grow horticultural crops throughout the year in Haryana, production is restricted because of water deficit. Rainfall is insufficient to maintain adequate soil moisture levels for crop production particularly during the dry season (December to June). Consequently, crops experience water stress and irrigation needs to be ensured to obtain acceptable yields. Existing ponds and dams in Haryana are insufficient to store water for agricultural purposes and the low quality of underground water restricts production of many horticultural crops. Rainwater catchments have long been used in Haryana for domestic consumption and appear to be a feasible supply of good quality water for high value horticultural crops. Adoption of ground liners as a rainwater catchments system has been

difficult because it requires additional land that is lost for production and the system has been focused to irrigate crops under full sun that require high amount of water.

REVIEW OF LITERATURE

The water deficit in Haryana horticultural industry can be ameliorated by growing crops under shade since plants require substantially less irrigation water. Crops grown in open fields of a semi-dry tropical climate are subjected to direct sunlight, high temperatures and wind resulting in high crop evapo-transpiration (ET_c) and therefore, demanding large amounts of water. In contrast, shade-houses favor plant growth since plants are less stressful: direct sunlight is avoided, temperature is lower, humidity is higher, wind is reduced, and ET_c is low. Irrigation water requirement of 23% to 31% pan evaporation has been used for leatherleaf fern grown under 70% light reduction (Stamps, 1995). In addition, water use efficiency increases under shady conditions (Durr and Rangel, 2003;

Jifon and Syvertsen, 2003). Excessive shade, however, can be detrimental for crop production because of reduced photosynthesis. This suggests that Haryana horticultural industry may benefit tremendously by the incorporation of shade-houses for a water sustainable production system. In addition, some high value ornamentals perform better when grown under shady conditions and there is a demand to complement the flourishing tourist industry in Haryana.

This study evaluated water requirements to produce anthurium cut flowers and heliconias under shade conditions in Haryana. The specific objectives of this study are: a) to determine the water use efficiency in crops grown under shade conditions, and b) to determine the optimal shade for production of anthurium and heliconias. Crops were grown in a shade-house with three different shade levels and the irrigation water balance was analyzed and compared with meteorological data. The information obtained in this project will help horticulturists in Haryana develop decision criteria for crop selection and adoption of shade-houses to improve water productivity (WP).

METHODOLOGY AND PRINCIPAL FINDINGS

Experiment I

Methodology:

In the first experiment, three water stress levels and two shade levels were studied for Anthurium cut flower production. Irrigation schedule was based on soil moisture levels at -10, -30, and -50 kPa to determine the effect of water stress on water savings and flower production and quality. The shade levels were 60% and 80%. Eight plants were planted on 1.44 m² beds filled with a base of 20 cm sand topped with another

20 cm soilless media (Promix BX). Three plots (repetitions) were set up for each treatment. Plots were irrigated with micro-sprinklers scheduled to irrigate when soil moisture reached the corresponding treatment level. Soil moisture was monitored by Watermark soil moisture sensors (Irrometer Co., Riverside, CA) and connected to the Watermark Electronic Module (Irrometer Co.) which would allow the solenoid to open for irrigation. Soil moisture, applied water quantity, pan-evaporation, and flower production were monitored from January to April, 2006 to determine water requirements. Rainfall and pan evaporation from the meteorological station was monitored also. Due to the inconsistency of the Watermark sensor at stressing levels and the low quality of the flowers, the irrigation schedule was changed in May to daily irrigation with fixed amount of water.

Treatment consisted in 1.1, 2.2 and 3.3 mm a day for both shade levels. These amounts were approximately 16%, 33% and 50% pan evaporation at full sun. In addition, 2.2 and 3.3 mm correspond to 1.2 times pan evaporation at 80% and 60% shade, respectively.

Principal Findings:

Rainfall was substantially lower than evapo-transpiration particularly during the dry season (January to August) (table 1, fig 1). By the end of April when irrigation scheduling was changed, pan evaporation at 60% and 80% shade was reduced to 47% and 32% of full sun pan-evaporation, respectively, and rainfall was 21% (79% deficit).

Similar differences in pan evaporation continued until the end of the experiments in November, however, rainfall deficit was reduced to 53% since the rain was more frequent and heavier. These results suggest that shading is an excellent alternative to reduce crop water requirements in comparison to direct sun production and that it is possible to grow crops in a shade-house modified to catch rainwater and use it for irrigation. In the first period, the automatic irrigation system worked well at -10 kPa allowing irrigation to start.

The amount of water applied to the -10kPa treatment at 60% and 80% shade was 66 mm in 4 applications and 33 mm in 2 applications, respectively. In contrast, at lower soil moisture levels the moisture sensors appeared to have lost contact with the media and reading became erratic. In this production period, the number of harvested flowers was slightly lower at 80% shade (fig 2), but flower quality appeared to be superior. In addition, water stress with the purpose of saving water was detrimental on flower production since the number of flowers (and quality) was dramatically reduced (fig 2). In the second production period, there was no difference in production between the amounts of applied water at 80% shade (fig 3). Yield of flowers was higher at

60% than at 80% shade suggesting that 80% shade may have been below the optimum and photosynthesis may have been reduced. Within 60% shade yield was reduced with lower amount of water suggesting that irrigation may have been suboptimal. Therefore, evapotranspiration needs to be satisfied in full for optimum production.

METHODOLOGY

A second experiment was conducted in September – November, 2006 to estimate water requirements and water use efficiency in production of potted anthurium and heliconia under three shade conditions. A greenhouse 20' by 12' was built and separated in four sections with shade levels of 0%, 30%, 50% and 70% (DeWitt Knitted shade cloth, Hummert International, Earth City, MO). Six anthuriums and three heliconias growing in three-gallon black pots were used per treatment. Rainwater collected from the greenhouse's top through gutters was stored and used for drip-irrigation and plants were fertilized by injecting soluble fertilizer (Peters 20-20-20 + micronutrients at 100ppm N).

An automatic irrigation systems based on a timer was set up to irrigate pots for 1 min three times a day. Drippers of 2, 1, 0.5, and 0.5gal.h⁻¹ were used for 0%, 30%, 50% and 70% shade, respectively. Excess drain water was monitored daily to determine daily water used and pot weigh was monitored once a month. Pan evaporation was monitored at each shade level for comparison with rainfall and pan evaporation from the meteorological station. Temperature, relative humidity and solar radiation were monitored also with a Watchdog mini-datalogger (Spectrum Technologies, Inc., East Plainfield, IL).

Principal Findings:

Water requirements of potted anthurium and heliconia were reduced significantly as shade level was increased from section 1 (no shade) to section 4 (70% shade) (table 2).

Similar to pan-evaporation, daily water used by both anthuriums and heliconias was reduced to 21% and 18%, respectively, from no shade to 70% shade. This reduction appears to be the effect of less solar radiation since temperature and relative humidity were similar among the shade levels. Pan evaporation recorded at the shade-house including the non-shaded treatment was lower than pan evaporation at the weather station. 55% reduction in section 1 which was not shaded may have been due to the reduced solar radiation under the plastic top and the more protected conditions that may have reduced the wind effect. Growth of the potted plants was not detected in the shaded treatments. Both anthurium and heliconias did not survive the non-shaded treatment and plants

turned yellow and 60 % died during the trial. Three month trial appears to be insufficient to evaluate growth of tropical ornamental under shade conditions.

Therefore, a longer testing period is needed to determine water productivity for these crop species.

Table 1

Total Evaporation and Rainfall, St Croix, USVI. January-November, 2006 Cumulative Percent from pan evaporation

	Cumulative pan evaporation	Rainfall	Evap I	Evap II	Percent from Pan evaporation		
					Rainfall	Evap I	Evap II
Shade	(Weather station		80%	60%		80%	60%
	mm	mm	mm	mm	%	%	%
July 2011	629	135	203	297	21	32	47
August 2011	1824	970	516	862	53	28	47

Cumulative Rainfall and Pan-evaporation at Full Sun, 60% and 80% Shade.

Table 2

Pan-evaporation, water used, temperature (temp), relative humidity (RH), and solar radiation (SRD) in shade-house production of potted anthuriums and heliconias

	Daily pan evaporation	Daily water used /plant				
	mm	Anthurium mL	Heliconia mL	temp C	RH %	SRD MJ/m2/day
Weather station	5.8					
0% shade	3.2	297a	265 a	28.2	77	14.6
30% shade	2.3	150 b	124 b	28.1	78	8.1
50% shade	2.0	75 c	68 c	28.1	78	6.3
70% shade	1.4	62 c	48 c	27.7	81	3.8

CONCLUSIONS

It may be concluded then that:

- Catching rainwater in a modified shade-house and use it for irrigation may be sufficient to implement a water sustainable production system.
- Increasing shade level reduces water requirements for Anthurium and heliconia production and yield. Anthurium and heliconia are recommended to be grown under 50% - 70% shade to improve growth and the flower quality.
- Stressing plants to save water is not recommended due to reduction of flower production and quality.

Development of a water sustainable production system will impact favorably the horticulture industry since water is the most limiting constraint to agriculture in the USVI. Water deficit ranges between 40% and 70% during the dry season in the territory.

Production of shade loving crops appears to be an excellent alternative to reduce water requirement and improve water use efficiency and productivity in Haryana. In addition, collection and storage of rainwater will reduce the dependency on costly city water. Rainwater collection and storage has being long adopted in Haryana for household consumption and can be easily adopted to develop a water sustainable production system.

Adoption of this system by small-scale growers may increase production of ornamental and specialty crops in Haryana impacting tremendously the local economy.

REFERENCES

1. Anis, M. and Iqbal, M. 1994. Medicinal plantlore of Aligarh, India. *Int. J. Pharmacog.* 32 (1): 59-64.
2. Prajapati, N.D., Purohit, S.S., Sharma, A.K. and Kumar, T. 2007. *A Handbook of Medicinal Plants*. Jodhpur, Agrobios (India), Section-II, pp. 1-554.
3. Jain, S.K. 1991. *Dictionary of Indian Folk Medicine and Ethnobotany*. New Delhi, Deep Publications, pp. 1-311.
4. Jain, S.R. and Sharma, S.N. 1967. Hypoglycaemic drugs of Indian indigenous origin. *Planta. Med.* 15 (4): 439-442.
5. Himalaya healthcare products, <http://www.himalayahealthcare.com/products.htm> downloaded on July 2, 2008.
6. Behl, H.M., Sidhu, O.P., Mehrotra, S., Pushpangadan, P. and Singh, S.C. 2006. *Nontoxic dental care herbal formulation for preventing dental plaque and gingivitis*. CSIR, New Delhi. (United States Patent No.7,083,779; <http://patft.uspto.gov>).

7. Reddy, M.B., Reddy, K.R. and Reddy, M.N. 1988. A Survey of medicinal plants of Chenchu tribes of Andhra Pradesh, India. *Int. J. Crude Drug Res.* 26 (4): 189-196.

8. Arseculeratne, S.N., Gunatilaka, A.A.L. and Panabokke, R.G. 1985. Studies on medicinal plants of Sri Lanka. Part 14. Toxicity of some traditional medicinal herbs. *J. Ethnopharmacol.* 13 (3): 323-335.

9. Gaitonde, B.B., Kulharni, H.J., Nabar, S.D. and Joglekar, S.N. 1974. Diuretic activity of Punarnava (*Boerhavia diffusa*). *Bull. Haffkine Inst.* (2): p. 24.

10. Olukoya, D.K., Idika, N. and Odugbemi, T. 1993. Antibacterial activity of some medicinal plants from Nigeria. *J. Ethnopharmacol.* 39 (1): 69-72.

11. Organic ayurvedic herbal products, <http://www.ayurveda.cz/en/organic-ayurvedic-herbal-products.htm> downloaded on June 30, 2008.

12. Bapurao, M.N. 2005. Herbal composition for the treatment of hepatic and splenic disorders. *Sahajanand Biotech Private Ltd (India), (Publication number & date*

GB2404147, 2005-01-26; http://v3.espacenet.com).

13. Tomer, O.S., Glomski, P. and Borah, K. 2000. Herbal compositions and their use as agents for control of hypertension, hypercholesterolemia and hyperlipidemia. *New Jersey,*

Chromak Research Inc. (United States Patent No. 6,162,438; http://patft.uspto.gov).

14. Duke, J.A. and Vasquez, R. 1994. *Amazonian Ethnobotanical Dictionary*. Boca Raton, CRC Press, p. 181.

15. Coee, F.G. and Anderson, G.J. 1996. Ethnobotany of the Garifuna of Eastern Nicaragua. *Econ. Bot.* 50 (1): 71-107.

16. Gupta, M.P., Arias, T.D., Correa, M. and Lamba, S.S. 1979. Ethnopharmacognosic observations on Panamanian medicinal plants. Part I. *Q. J. Crude Drug Res.* 17 (3/4): 115-

130.

17. Wild crafted herbal products, <http://www.wildcrafted.com.au/Moisturisers.html> downloaded on June 30, 2008.

18. Agarwal, R.K. and Agarwal, A. 2004. *Herbal composition having antiallergic properties and a process for the preparation thereof*. Bangalore, Natural Remedies Pvt. Ltd.

(United States Patent No. 6,730,332;
<http://patft.uspto.gov>).

19. Morimoto, C. and Dang, N.H. 2006. *Compositions for cancer prevention, treatment, or amelioration comprising papaya extract*. (Publication number & date WO/2006/

004226, 12.01.2006;
<http://www.wipo.int/pctdb/en/wo.jsp>).

20. Reddy, M.B., Reddy, K.R. and Reddy, M.N. 1989. A survey of plant crude drugs of Anantapur district, Andhra Pradesh, India. *Int. J. Crude Drug Res.* 27 (3): 145-155.

21. Bellakhdar, J., Claisse, R., Fleurentin, J. and Younos, C. 1991. Repertory of standard herbal drugs in the Moroccan pharmacopoea. *J. Ethnopharmacol.* 35 (2): 123-143.

22. Al-khalil, S. 1995. A Survey of plants used in Jordanian traditional medicine. *Int. J. Pharmacog.* 33 (4): 317-323.

23. Beneforce-Holistic Medicine, Alternative Health and Complementary Alternative Medicine, <http://beneforce.com/informationfaq/homeopathic/citrus-colocynthis.htm> downloaded on June 29, 2008.

24. Manandhar, N. P. 1995. An inventory of some herbal drugs of Myagdi district, Nepal. *Econ. Bot.* 49 (4): 371-379.

25. Sahu, T. R. 1984. Less known uses of weeds as medicinal plants. *Ancient. Sci. Life.* 3 (4): 245-249.

26. Manandhar, N. P. 1995. An inventory of some vegetable drug resources of Makawanpur district, Nepal. *Fitoterapia* 66 (3): 231-238.

27. Tropilab, <http://tropilab.com/jatrophatincture.html> downloaded on June 30, 2008.

28. Pushpangadan, P., Mehrotra, S., Rawat, A.K.S., Rao, Ch.V., Ojha, S.K. and Aziz, I. 2008. *Herbal composition for cuts, burns and wounds*. CSIR, New Delhi. (United States